



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 413/10, A61K 31/42, C07D 263/24, 263/18, 417/10 // (C07D 413/10, 309:00) (C07D 413/10, 211:00)</p>	<p>A1</p>	<p>(11) International Publication Number: WO 97/30995</p> <p>(43) International Publication Date: 28 August 1997 (28.08.97)</p>
<p>(21) International Application Number: PCT/GB97/00462</p> <p>(22) International Filing Date: 20 February 1997 (20.02.97)</p> <p>(30) Priority Data: 9603939.1 24 February 1996 (24.02.96) GB 9618404.9 4 September 1996 (04.09.96) GB</p> <p>(71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB).</p> <p>(72) Inventor; and (75) Inventor/Applicant (for US only): GRAVESTOCK, Michael, Barry [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).</p> <p>(74) Agent: DENERLEY, Paul, Millington; Intellectual Property Dept., Zeneca Pharmaceuticals, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. With amended claims.</p>
<p>(54) Title: ANTIBIOTIC OXAZOLIDINONE DERIVATIVES</p> <div data-bbox="535 1113 1153 1365" data-label="Chemical-Block"> <p style="text-align: right;">(I)</p> </div> <p>(57) Abstract</p> <p>The invention concerns a compound of formula (I) wherein, for example: R¹ is of the formula -NHC(=O)R^b wherein R^b is, for example, (1-4C)alkyl; R² and R³ are hydrogen or fluoro; D is O, S, SO, SO₂ or NR⁷; R⁴ and R⁵ are hydrogen, AR-oxymethyl; R⁶ is hydrogen; >A-B- is of the formula >C=C(R^a)-, >CHCHR^a-, or >C(OH)CHR^a-, (>represents two single bonds) wherein R^a is hydrogen or (1-4C)alkyl; R⁷ is hydrogen, AR or of the formula R¹⁰CO-, R¹⁰SO₂-, R¹⁰CS-, R^dOC(R^e)=CH(C=O)-, R^fC(=O)C(=O)-, R^gN-C(R^h)C(=O)- or RⁱNHC(R^j)=CHC(=O)- or R¹⁴CH(R¹³)(CH₂)_m-; wherein, for example, R¹⁰ is AR; R^d, R^e, R^f, R^g, R^h, Rⁱ and R^j are (1-6C)alkyl; m is 0 or 1, R¹³ is cyano, R¹⁴ is hydrogen or (1-4C)alkyl; AR is phenyl; and pharmaceutically-acceptable salts thereof; processes for their preparation; pharmaceutical compositions containing them and their use as antibacterial agents.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

ANTIBIOTIC OXAZOLIDINONE DERIVATIVES

The present invention relates to antibiotic compounds and in particular to antibiotic compounds containing an oxazolidinone ring. This invention further relates to processes for
5 their preparation, to intermediates useful in their preparation, to their use as therapeutic agents and to pharmaceutical compositions containing them.

The international microbiological community continues to express serious concern that the evolution of antibiotic resistance could result in strains against which currently available antibacterial agents will be ineffective. In general, bacterial pathogens may be
10 classified as either Gram-positive or Gram-negative pathogens. Antibiotic compounds with effective activity against both Gram-positive and Gram-negative pathogens are generally regarded as having a broad spectrum of activity. The compounds of the present invention are regarded primarily as effective against Gram-positive pathogens because of their particularly good activity against such pathogens.


15 Gram-positive pathogens, for example Staphylococci, Enterococci, Streptococci and mycobacteria, are particularly important because of the development of resistant strains which are both difficult to treat and difficult to eradicate from the hospital environment once established. Examples of such strains are methicillin resistant staphylococcus (MRSA), methicillin resistant coagulase negative staphylococci (MRCNS), penicillin resistant
20 streptococcus pneumoniae and multiply resistant Enterococcus faecium.

The major clinically effective antibiotic for treatment of such resistant Gram-positive pathogens is vancomycin. Vancomycin is a glycopeptide and is associated with nephrotoxicity and ototoxicity. Furthermore, and most importantly, antibacterial resistance to vancomycin and other glycopeptides is also appearing. This resistance is
25 increasing at a steady rate rendering these agents less and less effective in the treatment of Gram-positive pathogens.

The present inventors have discovered a class of antibiotic compounds containing an oxazolidinone ring which has useful activity against Gram-positive pathogens including MRSA and MRCNS and, in particular, against various strains exhibiting resistance to
30 vancomycin and against E. faecium strains resistant to both aminoglycosides and clinically used β -lactams.

We have now discovered a narrow range of compounds that is not suggested by the art and which has good activity against a broad range of Gram-positive pathogens including organisms known to be resistant to most commonly used antibiotics. In comparison with compounds described in the art (for example Walter A. Gregory et al in J.Med.Chem. 1990, 33, 2569-2578 and Chung-Ho Park et al in J.Med.Chem. 1992, 35, 1156-1165) the compounds also possess a favourable toxicological profile.

Accordingly the present invention provides a compound of the formula (I):



10

(I)

wherein:

- R¹ is hydroxy, chloro, fluoro, (1-4C)alkanesulfonyloxy, amino, azido, (1-4C)alkoxy,
 15 (1-4C)alkylthio, (1-4C)alkylaminocarbonyloxy;
 or of the formula -NHC(=O)R^b wherein R^b is hydrogen, (1-4C)alkoxy, amino, chloromethyl, dichloromethyl, cyanomethyl, methoxymethyl, acetylmethyl, methylamino, dimethylamino or (1-4C)alkyl;
 or R¹ is of the formula -NHS(O)_n(1-4C)alkyl wherein n is 0, 1 or 2;
 20 R² and R³ are independently hydrogen or fluoro;
 D is O, S, SO, SO₂ or NR⁷;
 when D is O, R⁴ and R⁵ are independently hydroxy, bromo, oxo (=O), (1-4C)alkyl, (1-4C)alkanoylamino-(1-4C)alkyl, hydroxy-(1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl, AR-oxymethyl, AR-thiomethyl (wherein AR is as defined hereinbelow) or independently as
 25 defined for R⁷ hereinbelow;
 when D is S, SO, SO₂ or NR⁷, R⁴ and R⁵ are independently oxo (=O), (1-4C)alkyl, (1-4C)alkanoylamino-(1-4C)alkyl, hydroxy-(1-4C)alkyl, carboxy,

(1-4C)alkoxycarbonyl, AR-oxymethyl, AR-thiomethyl (wherein AR is as defined hereinbelow) or independently as defined for R' hereinbelow;

R⁶ is hydrogen, (1-4C)alkyl, hydroxy, (1-4C)alkoxy or (2-4C)alkanoyloxy;

>A-B- is of the formula >C=C(R^a)-, >CHCHR^a- or >C(OH)CHR^a- (> represents two single bonds) wherein R^a is hydrogen or (1-4C)alkyl;

R' is hydrogen, cyano, 2-((1-4C)alkoxycarbonyl)ethenyl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl, 2-((1-4C)alkylaminocarbonyl)ethenyl, AR (as defined hereinbelow) or a tetrazole ring system (optionally mono-substituted in the 1- or 2- position of the tetrazole ring) wherein the tetrazole ring system is joined to the nitrogen in NR' by a ring carbon atom;

or R' is of the formula R¹⁰CO-, R¹⁰SO₂- or R¹⁰CS- wherein R¹⁰ is AR (as defined hereinbelow), cyclopentyl or cyclohexyl (wherein the last two-mentioned cycloalkyl rings are optionally mono- or disubstituted by substituents independently selected from (1-4C)alkyl (including geminal disubstitution), hydroxy,

(1-4C)alkoxy, (1-4C)alkylthio, acetamido, (1-4C)alkanoyl, cyano and trifluoromethyl), (1-4C)alkoxycarbonyl, hydrogen, amino, trifluoromethyl, (1-4C)alkylamino, di((1-4C)alkyl)amino, 2,3-dihydro-5-oxothiazolo-[3,2-A]pyrimidin-6-yl, 2-(2-furyl)ethenyl, 2-(2-thienyl)ethenyl, 2-phenylethenyl (wherein the phenyl substituent is optionally substituted by up to three substituents independently selected from (1-4C)alkoxy, halo and cyano), 3,4-dihydropyran-2-yl, coumal-5-yl, 5-methoxy-4-oxopyran-2-yl, N-acetylpyrrolidin-2-yl, 5-oxo-tetrahydrofuran-2-yl, benzopyranone or (1-10C)alkyl [wherein (1-10C)alkyl is optionally substituted by hydroxy, cyano, halo, (1-10C)alkoxy, trifluoromethyl, (1-4C)alkoxy-(1-4C)alkoxy, (1-4C)alkoxy-(1-4C)alkoxy-(1-4C)alkoxy, (1-4C)alkanoyl, (1-4C)alkoxycarbonyl, amino, (1-4C)alkylamino, di((1-4C)alkyl)amino, (1-6C)alkanoylamino, (1-4C)alkoxycarbonylamino, N-(1-4C)alkyl-N-(2-6C)alkanoylamino, (1-4C)alkylS(O)_pNH-, (1-4C)alkylS(O)_p((1-4C)alkyl)N-, fluoro(1-4C)alkylS(O)_pNH-, fluoro(1-4C)alkylS(O)_p((1-4C)alkyl)N-, phosphono, (1-4C)alkoxy(hydroxy)phosphoryl, di-(1-4C)alkoxyphosphoryl, (1-4C)alkylS(O)_q-, phenylS(O)_q- (wherein the phenyl group is optionally substituted by up to three substituents independently selected from (1-4C)alkoxy, halo and cyano), or CY (as defined hereinbelow), wherein p is 1 or 2 and q is 0, 1 or 2];

- or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is an optionally substituted 5- or 6-membered heteroaryl, optionally substituted phenyl, (1-4C)alkylamino, benzyloxy-(1-4C)alkyl or optionally substituted (1-10C)alkyl;
- or R^{10} is of the formula $R^{12}O-$ wherein R^{12} is optionally substituted (1-6C)alkyl;
- 5 or R^7 is of the formula $R^dOC(R^e)=CH(C=O)-$, $R^fC(=O)C(=O)-$, $R^gN=C(R^h)C(=O)-$ or $R^iNHC(R^j)=CHC(=O)-$ wherein R^d is (1-6C)alkyl, R^e is hydrogen or (1-6C)alkyl, or R^d and R^e together form a (3-4C)alkylene chain, R^f is hydrogen, (1-6C)alkyl, hydroxy(1-6C)alkyl, (1-6C)alkoxy(1-6C)alkyl, amino, (1-4C)alkylamino, di-(1-4C)alkylamino, (1-6C)alkoxy, (1-6C)alkoxy(1-6C)alkoxy,
- 10 hydroxy(2-6C)alkoxy, (1-4C)alkylamino(2-6C)alkoxy, di-(1-4C)alkylamino(2-6C)alkoxy, R^g is (1-6C)alkyl, hydroxy or (1-6C)alkoxy, R^h is hydrogen or (1-6C)alkyl, R^i is hydrogen, (1-6C)alkyl, optionally substituted phenyl or an optionally substituted 5- or 6-membered heteroaryl and R^j is hydrogen or (1-6C)alkyl;
- or R^7 is of the formula $R^{14}CH(R^{13})(CH_2)_m-$ wherein m is 0 or 1, R^{13} is fluoro, cyano,
- 15 (1-4C)alkoxy, (1-4C)alkylsulfonyl, (1-4C)alkoxycarbonyl or hydroxy, (provided that when m is 0, R^{13} is not fluoro or hydroxy) and R^{14} is hydrogen or (1-4C)alkyl;
- wherein AR is optionally substituted phenyl, optionally substituted phenyl(1-4C)alkyl, optionally substituted 5- or 6-membered heteroaryl, optionally substituted naphthyl or an optionally substituted 5/6 or 6/6 bicyclic heteroaryl ring system, in which the bicyclic
- 20 heteroaryl ring systems may be linked via an atom in either of the rings comprising the bicyclic system, and wherein the mono- and bicyclic heteroaryl ring systems are linked via a ring carbon atom;
- wherein CY is a 4-, 5- or 6-membered cycloalkyl ring, a 5- or 6-membered cycloalkenyl ring, naphthoxy, thiophen-2-yl, indol-1-yl, indol-3-yl, pyrimidin-2-ylthio,
- 25 1,4-benzodioxan-6-yl, sulfolan-3-yl, pyridin-2-yl; wherein any of the afore-mentioned ring systems in CY may be optionally substituted by up to three substituents independently selected from halo, (1-4C)alkyl (including geminal disubstitution when CY is a cycloalkyl or cycloalkenyl ring), acyl, oxo and nitro-(1-4C)alkyl; and pharmaceutically-acceptable salts thereof; except that
- 30 N-((5S)-3-(4-(2-oxo-5,6-dihydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide is excluded.

In this specification a '5- or 6-membered heteroaryl' and 'heteroaryl (monocyclic) ring' means a 5- or 6-membered aryl ring wherein 1, 2 or 3 of the ring atoms are selected from nitrogen, oxygen and sulfur. Particular examples of 5- or 6-membered heteroaryl ring systems are furan, pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole and thiophene.

In this specification a '5/6 or 6/6 bicyclic heteroaryl ring system' and 'heteroaryl (bicyclic) ring' means an aromatic bicyclic ring system comprising a 6-membered ring fused to either a 5 membered ring or another 6 membered ring, the bicyclic ring system containing 10 to 4 heteroatoms selected from nitrogen, oxygen and sulfur. Particular examples of 5/6 and 6/6 bicyclic ring systems are indole, benzofuran, benzoimidazole, benzothiophene, benzisothiazole, benzoxazole, benzisoxazole, pyridoimidazole, pyrimidoimidazole, quinoline, quinoxaline, quinazoline, phthalazine, cinnoline and naphthyridine.

In this specification a '4-, 5- or 6-membered cycloalkyl ring' means a cyclobutyl, 15 cyclopentyl or cyclohexyl ring; and a '5- or 6-membered cycloalkenyl ring' means a cyclopentenyl or cyclohexenyl ring.

In this specification the term 'alkyl' includes straight chained and branched structures. For example, (1-6C)alkyl includes propyl, isopropyl and tert-butyl. However, references to individual alkyl groups such as "propyl" are specific for the straight chained 20 version only, and references to individual branched chain alkyl groups such as "isopropyl" are specific for the branched chain version only. A similar convention applies to other radicals, for example halo(1-4C)alkyl includes 1-bromoethyl and 2-bromoethyl.

Particular optional substituents for alkyl, phenyl (and phenyl containing moieties) and naphthyl groups and ring carbon atoms in heteroaryl (mono or bicyclic) rings in R¹¹, R¹², 25 Rⁱ and AR include halo, (1-4C)alkyl, hydroxy, nitro, carbamoyl, (1-4C)alkylcarbamoyl, di-((1-4C)alkyl)carbamoyl, cyano, trifluoromethyl, trifluoromethoxy, amino, (1-4C)alkylamino, di((1-4C)alkyl)amino, (1-4C)alkyl S(O)_q, (wherein q is 0, 1 or 2), carboxy, (1-4C)alkoxycarbonyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkanoyl, (1-4C)alkoxy, (1-4C)alkanoylamino, benzoylamino, benzoyl, phenyl (optionally substituted by up to three 30 substituents selected from halo, (1-4C)alkoxy or cyano), furan, pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole, thiophene.

hydroxyimino(1-4C)alkyl, (1-4C)alkoxyimino(1-4C)alkyl, hydroxy-(1-4C)alkyl, halo-(1-4C)alkyl, nitro(1-4C)alkyl, amino(1-4C)alkyl, cyano(1-4C)alkyl, (1-4C)alkanesulfonamido, aminosulfonyl, (1-4C)alkylaminosulfonyl and di-((1-4C)alkyl)aminosulfonyl. The phenyl and naphthyl groups and heteroaryl (mono- or bicyclic) rings in R¹¹, Rⁱ and AR may be mono-
 5 or disubstituted on ring carbon atoms with substituents independently selected from the above list of particular optional substituents.

Particular optional substituents for ring nitrogen atoms when R⁷ is tetrazole, in heteroaryl groups in R¹¹, R¹², Rⁱ and AR, and in the nitrogen-containing rings in CY, which can be substituted without becoming quaternised include (1-4C)alkyl, (2-4C)alkenyl, (2-
 10 4C)alkynyl and (1-4C)alkanoyl.

Examples of halo groups include fluoro, chloro and bromo; examples of (1-4C)alkyl, include methyl, ethyl, and propyl and isopropyl; examples of (1-6C)alkyl include methyl, ethyl, propyl, isopropyl, pentyl and hexyl; examples of (1-10C)alkyl include methyl, ethyl, propyl, isopropyl, pentyl, hexyl, heptyl, octyl and nonyl; examples of (1-4C)alkylamino
 15 include methylamino, ethylamino and propylamino; examples of di-((1-4C)alkyl)amino include dimethylamino, N-ethyl-N-methylamino, diethylamino, N-methyl-N-propylamino and dipropylamino; examples of (1-4C)alkylS(O)_q wherein q is 0, 1 or 2 include methylthio, ethylthio, methylsulfinyl, ethylsulfinyl, methylsulfonyl and ethylsulfonyl; examples of (1-4C)alkanesulfonyloxy include methylsulfonyloxy, ethylsulfonyloxy and
 20 propylsulfonyloxy; examples of (1-4C)alkylthio include methylthio and ethylthio; examples of (1-4C)alkylsulfonyl include methylsulfonyl and ethylsulfonyl; examples of (1-4C)alkylaminocarbonyloxy include methylaminocarbonyloxy and ethylaminocarbonyloxy; examples of (1-4C)alkanoylamino-(1-4C)alkyl include formamidomethyl, acetamidomethyl and acetamidoethyl; examples of (1-6C)alkoxy-(1-6C)alkyl include methoxymethyl,
 25 ethoxymethyl and 2-methoxyethyl; examples of (1-4C)alkoxycarbonyl include methoxycarbonyl, ethoxycarbonyl and propoxycarbonyl; examples of (2-4C)alkanoyloxy include acetyloxy and propionyloxy; examples of (1-4C)alkoxy include methoxy, ethoxy and propoxy; examples of (1-6C)alkoxy include methoxy, ethoxy, propoxy and pentoxy; examples of hydroxy-(2-6C)alkoxy include 2-hydroxyethoxy and 3-hydroxypropoxy; examples of
 30 (1-4C)alkylamino-(2-6C)alkoxy include 2-methylaminoethoxy and 2-ethylaminoethoxy; examples of di-(1-4C)alkylamino-(2-6C)alkoxy include 2-dimethylaminoethoxy and

2-diethylaminoethoxy; examples of (1-4C)alkoxy-(1-4C)alkoxy and (1-6C)alkoxy-(1-6C)alkoxy include methoxymethoxy, 2-methoxyethoxy, 2-ethoxyethoxy and 3-methoxypropoxy; examples of (1-4C)alkoxy-(1-4C)alkoxy-(1-4C)alkoxy include 2-(methoxymethoxy)ethoxy, 2-(2-methoxyethoxy)ethoxy; 3-(2-methoxyethoxy)propoxy and 5 2-(2-ethoxyethoxy)ethoxy; examples of (1-4C)alkanoylamino and (1-6C)alkanoylamino include formamido, acetamido and propionylamino; examples of (1-4C)alkoxycarbonylamino include methoxycarbonylamino and ethoxycarbonylamino; examples of N-(1-4C)alkyl-N-(2-6C)alkanoylamino include N-methylacetamido, N-ethylacetamido and N-methylpropionamido; examples of (1-4C)alkylS(O)_pNH- wherein p is 1 or 2 include 10 methylsulfinylamino, methylsulfonylamino, ethylsulfinylamino and ethylsulfonylamino; examples of (1-4C)alkylS(O)_p((1-4C)alkyl)NH- wherein p is 1 or 2 include methylsulfinylmethylamino, methylsulfonylmethylamino, 2-(ethylsulfinyl)ethylamino and 2-(ethylsulfonyl)ethylamino; examples of fluoro(1-4C)alkylS(O)_pNH- wherein p is 1 or 2 include trifluoromethylsulfinylamino and trifluoromethylsulfonylamino; examples of 15 fluoro(1-4C)alkylS(O)_p((1-4C)alkyl)NH- wherein p is 1 or 2 include trifluoromethylsulfinylmethylamino and trifluoromethylsulfonylmethylamino examples of (1-4C)alkoxy(hydroxy)phosphoryl include methoxy(hydroxy)phosphoryl and ethoxy(hydroxy)phosphoryl; examples of di-(1-4C)alkoxyphosphoryl include di-methoxyphosphoryl, di-ethoxyphosphoryl and ethoxy(methoxy)phosphoryl; examples of 20 2-((1-4C)alkoxycarbonyl)ethenyl include 2-(methoxycarbonyl)ethenyl and 2-(ethoxycarbonyl)ethenyl; examples of 2-cyano-2-((1-4C)alkyl)ethenyl include 2-cyano-2-methylethenyl and 2-cyano-2-ethylethenyl; examples of 2-((1-4C)alkylaminocarbonyl)ethenyl include 2-(methylaminocarbonyl)ethenyl and 2-(ethylaminocarbonyl)ethenyl; examples of benzyloxy(1-4C)alkyl include benzyloxymethyl 25 and benzyloxyethyl; examples of phenyl(1-4C)alkyl include benzyl and phenethyl; examples of phenylS(O)_q wherein q is 0, 1 or 2 are phenylthio, phenylsulfinyl and phenylsulfonyl respectively; examples of (1-4C)alkylcarbamoyl include methylcarbamoyl and ethylcarbamoyl; examples of di((1-4C)alkyl)carbamoyl include di(methyl)carbamoyl and di(ethyl)carbamoyl; examples of a (3-4C)alkylene chain are trimethylene or tetramethylene; 30 examples of (2-4C)alkenyl include allyl and vinyl; examples of (2-4C)alkynyl include ethynyl and

- 2-propynyl; examples of (1-4C)alkanoyl include formyl, acetyl and propionyl; examples of hydroxyimino(1-4C)alkyl include hydroxyiminomethyl, 2-(hydroxyimino)ethyl and 1-(hydroxyimino)ethyl; examples of (1-4C)alkoxyimino-(1-4C)alkyl include methoxyiminomethyl, ethoxyiminomethyl, 1-(methoxyimino)ethyl and 2-(methoxyimino)ethyl; examples of hydroxy(1-4C)alkyl and hydroxy(1-6C)alkyl include hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl and 3-hydroxypropyl; examples of halo(1-4C)alkyl include, halomethyl, 1-haloethyl, 2-haloethyl, and 3-halopropyl; examples of nitro(1-4C)alkyl include nitromethyl, 1-nitroethyl, 2-nitroethyl and 3-nitropropyl; examples of amino(1-4C)alkyl include aminomethyl, 1-aminoethyl, 2-aminoethyl and 3-aminopropyl; examples of cyano(1-4C)alkyl include cyanomethyl, 1-cyanoethyl, 2-cyanoethyl and 3-cyanopropyl; examples of (1-4C)alkanesulfonamido include methanesulfonamido and ethanesulfonamido; examples of (1-4C)alkylaminosulfonyl include methylaminosulfonyl and ethylaminosulfonyl; and examples of di-(1-4C)alkylaminosulfonyl include dimethylaminosulfonyl, diethylaminosulfonyl and N-methyl-N-ethylaminosulfonyl.
- 15 Suitable pharmaceutically-acceptable salts include acid addition salts such as methanesulfonate, fumarate, hydrochloride, hydrobromide, citrate, maleate and salts formed with phosphoric and sulfuric acid. In another aspect suitable salts are base salts such as an alkali metal salt for example sodium, an alkaline earth metal salt for example calcium or magnesium, an organic amine salt for example triethylamine, morpholine, N-methylpiperidine, N-ethylpiperidine, procaine, dibenzylamine, N,N-dibenzylethylamine or amino acids for example lysine. There may be more than one cation or anion depending on the number of charged functions and the valency of the cations or anions. A preferred pharmaceutically-acceptable salt is the sodium salt.

25 However, to facilitate isolation of the salt during preparation, salts which are less soluble in the chosen solvent may be preferred whether pharmaceutically-acceptable or not.


 The compounds of the formula (I) may be administered in the form of a pro-drug which is broken down in the human or animal body to give a compound of the formula (I). Examples of pro-drugs include in-vivo hydrolysable esters of a compound of the formula (I).

30 An in-vivo hydrolysable ester of a compound of the formula (I) containing carboxy or hydroxy group is, for example, a pharmaceutically-acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically-

acceptable esters for carboxy include (1-6C)alkoxymethyl esters for example methoxymethyl, (1-6C)alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, (3-8C)cycloalkoxycarbonyloxy(1-6C)alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and 5 (1-6C)alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be formed at any carboxy group in the compounds of this invention.

An in-vivo hydrolysable ester of a compound of the formula (I) containing a hydroxy group includes inorganic esters such as phosphate esters and α -acyloxyalkyl ethers and related compounds which as a result of the in-vivo hydrolysis of the ester breakdown to 10 give the parent hydroxy group. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxymethoxy. A selection of in-vivo hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and N-(dialkylaminoethyl)-N-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and 15 carboxyacetyl.

The compounds of the present invention have a chiral centre at the C-5 position of the oxazolidinone ring. The pharmaceutically active enantiomer is of the formula:



20

(IA)

The present invention includes the pure enantiomer depicted above or mixtures of the 5R and 5S enantiomers, for example a racemic mixture. If a mixture of enantiomers is 25 used, a larger amount (depending upon the ratio of the enantiomers) will be required to achieve the same effect as the same weight of the pharmaceutically active enantiomer. For the avoidance of doubt the enantiomer depicted above could be either 5R or 5S depending upon

the value of R^1 . For example, when R^1 is acetamido, the enantiomer depicted above is the 5S enantiomer and when R^1 is hydroxy, the enantiomer depicted above is the 5R enantiomer.

Furthermore, some compounds of the formula (I) may have other chiral centres, and some compounds of the formula (I) may exist as one or more regioisomers. It is to be understood that the invention encompasses all such optical, diastereo- and regio-isomers that possess antibacterial activity.

The invention relates to all tautomeric forms of the compounds of the formula (I) that possess antibacterial activity.

It is also to be understood that certain compounds of the formula (I) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess antibacterial activity.

In a preferred aspect of the invention there is provided a compound of the formula (I) as defined above, except that in the definition of R^4 and R^5 , neither R^4 nor R^5 can be oxo ($=O$).

In another aspect the present invention provides a compound of the formula (I) wherein:

R^1 is hydroxy, chloro, fluoro, (1-4C)alkanesulfonyloxy, amino, azido, (1-4C)alkoxy, (1-4C)alkylthio, (1-4C)alkylaminocarbonyloxy, or of the formula $-NHC(=O)R^b$ wherein R^b is hydrogen, (1-4C)alkoxy, amino, chloromethyl, dichloromethyl, cyanomethyl, methoxymethyl, acetylmethyl, methylamino, dimethylamino or (1-4C)alkyl;

or R^1 is of the formula $-NHS(O)_n(1-4C)alkyl$ wherein n is 0, 1 or 2;

R^2 and R^3 are independently hydrogen or fluoro;

when D is O, R^4 and R^5 are independently hydrogen, hydroxy, bromo, oxo ($=O$), (1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl or independently as defined for R^7 hereinbelow;

when D is S, SO, SO_2 or NR^7 , R^4 and R^5 are independently hydrogen, oxo ($=O$), (1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl or independently as defined for R^7 hereinbelow;

R^6 is hydrogen, (1-4C)alkyl, hydroxy, (1-4C)alkoxy or (2-4C)alkanoyloxy;

$>A-B-$ is of the formula $>C=C(R^a)$, $>CHCHR^a$, or $>C(OH)CHR^a$ ($>$ represents two single bonds) wherein R^a is hydrogen or (1-4C)alkyl;

D is O, S, SO, SO_2 or NR^7 ;

the value of R^1 . For example, when R^1 is acetamido, the enantiomer depicted above is the 5S enantiomer and when R^1 is hydroxy, the enantiomer depicted above is the 5R enantiomer.

Furthermore, some compounds of the formula (I) may have other chiral centres, and some compounds of the formula (I) may exist as one or more regioisomers. It is to be understood that the invention encompasses all such optical, diastereo- and regio-isomers that possess antibacterial activity.

The invention relates to all tautomeric forms of the compounds of the formula (I) that possess antibacterial activity.

It is also to be understood that certain compounds of the formula (I) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess antibacterial activity.

In a preferred aspect of the invention there is provided a compound of the formula (I) as defined above, except that in the definition of R^4 and R^5 , neither R^4 nor R^5 can be oxo (=O).

In another aspect the present invention provides a compound of the formula (I) wherein:

R^1 is hydroxy, chloro, fluoro, (1-4C)alkanesulfonyloxy, amino, azido, (1-4C)alkoxy, (1-4C)alkylthio, (1-4C)alkylaminocarbonyloxy, or of the formula $-NHC(=O)R^b$ wherein R^b is hydrogen, (1-4C)alkoxy, amino, chloromethyl, dichloromethyl, cyanomethyl, methoxymethyl, acetylmethyl, methylamino, dimethylamino or (1-4C)alkyl;

or R^1 is of the formula $-NHS(O)_n(1-4C)alkyl$ wherein n is 0, 1 or 2;

R^2 and R^3 are independently hydrogen or fluoro;

when D is O, R^4 and R^5 are independently hydrogen, hydroxy, bromo, oxo (=O), (1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl or independently as defined for R^7 hereinbelow;

when D is S, SO, SO_2 or NR^7 , R^4 and R^5 are independently hydrogen, oxo (=O), (1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl or independently as defined for R^7 hereinbelow;

R^6 is hydrogen, (1-4C)alkyl, hydroxy, (1-4C)alkoxy or (2-4C)alkanoyloxy;

$>A-B-$ is of the formula $>C=C(R^a)$, $>CHCHR^a$, or $>C(OH)CHR^a$ ($>$ represents two single bonds) wherein R^a is hydrogen or (1-4C)alkyl;

D is O, S, SO, SO_2 or NR^7 ;

- wherein R^7 is hydrogen, cyano, 2-((1-4C)alkoxycarbonyl)ethenyl, 2-((1-4C)alkylaminocarbonyl)ethenyl, optionally substituted phenyl, optionally substituted phenyl(1-4C)alkyl, optionally substituted 5- or 6-membered heteroaryl, optionally substituted naphthyl or optionally substituted 5/6 or 6/6 bicyclic heteroaryl ring system wherein the
- 5 heteroaryl ring systems are joined to the the nitrogen by a ring carbon atom;
or R^7 is of the formula $R^{10}CO-$ or $R^{10}SO_2-$
wherein R^{10} is (1-4C)alkoxycarbonyl, amino, (1-4C)alkylamino, di((1-4C)alkyl)amino or (1-6C)alkyl [wherein (1-6C)alkyl is optionally substituted by hydroxy, cyano, amino, (1-4C)alkoxy, (1-4C)alkanoyl, (1-4C)alkylamino, di((1-4C)alkyl)amino,
- 10 (2-6C)alkanoylamino, N-(1-4C)alkyl-N-(2-6C)alkanoylamino, (1-4C)alkylS(O)_pNH-, (1-4C)alkylS(O)_p((1-4C)alkyl)N-, phosphono, (1-4C)alkoxy(hydroxy)phoryl, di-(1-4C)alkoxyphosphoryl or (1-4C)alkylS(O)_q wherein p is 1 or 2 and q is 0, 1 or 2];
or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is optionally substituted 5- or 6-membered heteroaryl, optionally substituted phenyl or optionally substituted (1-6C)alkyl;
- 15 or R^{10} is of the formula $R^{12}O-$ wherein R^{12} is optionally substituted (1-6C)alkyl;
or R^7 is of the formula $R^dOC(R^e)=CH(C=O)-$, $R^fC(=O)C(=O)-$, $R^gN=C(R^h)C(=O)-$ or $R^iNHC(R^j)=CHC(=O)-$ wherein R^d is (1-6C)alkyl, R^e is hydrogen or (1-6C)alkyl or R^d and R^e together form a (3-4C)alkylene chain, R^f is hydrogen, (1-6C)alkyl, hydroxy(1-6C)alkyl, (1-6C)alkoxy(1-6C)alkyl, amino, (1-4C)alkylamino, di-(1-4C)alkylamino, (1-6C)alkoxy,
- 20 (1-6C)alkoxy(1-6C)alkoxy, hydroxy(2-6C)alkoxy, (1-4C)alkylamino(2-6C)alkoxy, di-(1-4C)alkylamino(2-6C)alkoxy, R^g is (1-6C)alkyl, hydroxy or (1-6C)alkyl, R^h is hydrogen or (1-6C)alkyl, R^i is (1-6C)alkyl, phenyl or a 5- or 6-membered heteroaryl and R^j is hydrogen or (1-6C)alkyl;
or R^7 is of the formula $R^{14}CH(R^{13})(CH_2)_m$ - wherein m is 0 or 1, R^{13} is fluoro, cyano,
- 25 (1-4C)alkoxy, (1-4C)alkylsulfonyl, (1-4C)alkoxycarbonyl or hydroxy; (provided that when m is 0, R^{13} is not fluoro or hydroxy) and R^{14} is hydrogen or (1-4C)alkyl;
and pharmaceutically-acceptable salts thereof; except that
N-((5S)-3-(4-(2-oxo-5,6-dihydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
N-((5S)-3-(3-ethyl-4-(2-oxo-5,6-dihydrothiapyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide and N-((5S)-3-(3-hydroxy-4-(2-oxo-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide are excluded.
- 30

In a preferred aspect of the invention there is provided a compound of the formula (I) as defined above in the section relating to another aspect of the present invention, except that R² is hydrogen (rather than hydrogen or (1-4C)alkyl).

In another preferred aspect of the invention there is provided a compound of the formula (I) as defined above in the section relating to another aspect of the present invention, except that in the definition of R⁴ and R⁵, neither R⁴ nor R⁵ can be oxo (=O).

In a further aspect the present invention provides a compound of the formula (I) wherein:

R¹ is of the formula -NHC(=O)(1-4C)alkyl or -NHS(O)_n(1-4C)alkyl wherein n is 0, 1 or 2;

10 R² and R³ are independently hydrogen or fluoro;

R⁴ and R⁵ are independently hydrogen, or methyl;

>A-B- is of the formula >C=CH, >CHCH₂, or >C(OH)CH₂- (> represents two single bonds)

wherein R⁴ is hydrogen or (1-4C)alkyl;

D is O, S, SO, SO₂ or NR⁷;

15 wherein R⁷ is hydrogen, optionally substituted phenyl, optionally substituted phenyl(1-4C)alkyl, optionally substituted 5- or 6-membered heteroaryl, optionally substituted naphthyl or optionally substituted 5/6 or 6/6 bicyclic heteroaryl ring system wherein the heteroaryl ring systems are joined to the the nitrogen in NR⁷ by a ring carbon atom; or R⁷ is of the formula R¹⁰CO- or R¹⁰SO₂-

20 wherein R¹⁰ is amino, (1-4C)alkylamino, di((1-4C)alkyl)amino or

(1-6C)alkyl [wherein (1-6C)alkyl is optionally substituted by hydroxy, cyano, amino, or (1-4C)alkylS(O)_q wherein q is 1 or 2];

or R¹⁰ is of the formula R¹¹C(O)O(1-6C)alkyl wherein R¹¹ is optionally substituted 5- or 6-membered heteroaryl, optionally substituted phenyl or optionally substituted (1-6C)alkyl;

25 or R¹⁰ is of the formula R¹²O- wherein R¹² is optionally substituted (1-6C)alkyl;

or R⁷ is of the formula R¹⁴CH(R¹³)(CH₂)_m - wherein m is 0 or 1, R¹³ is fluoro, cyano, (1-4C)alkoxy, (1-4C)alkylsulfonyl, (1-4C)alkoxycarbonyl or hydroxy; (provided that when m is 0, R¹³ is not fluoro or hydroxy) and R¹⁴ is hydrogen or (1-4C)alkyl; and pharmaceutically-acceptable salts thereof.

30 In the sections above relating to another aspect of the present invention, and to a further aspect of the present invention, '5- or 6-membered heteroaryl' means a 5- or 6-

membered aryl ring wherein 1, 2 or 3 of the ring atoms are selected from nitrogen, oxygen and sulfur. Particular examples of such 5- or 6-membered heteroaryl ring systems are imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole and thiophene.

5 Also, in the sections above relating to another aspect of the present invention, and to a further aspect of the present invention, a '5/6 or 6/6 bicyclic heteroaryl ring system' means an aromatic bicyclic ring system comprising a 6-membered ring fused to either a 5 membered ring or another 6 membered ring, the bicyclic ring system containing 1 to 4 heteroatoms selected from nitrogen, oxygen and sulfur. Particular examples of 5/6 and 6/6 bicyclic ring
10 systems are benzofuran, benzoimidazole, benzothiophene, benzisothiazole, benzoxazole, benzisoxazole, pyridoimidazole, pyrimidoimidazole, quinoline, quinoxaline, quinazoline, phthalazine, cinnoline and naphthyridine.

Particularly preferred compounds of the invention comprise a compound of the formula (I), or pharmaceutically-acceptable salts thereof, wherein the substituents A, B, D, R¹
15 to R¹⁴ and other optional substituents mentioned above have the values disclosed hereinbefore, or any of the following values :

(a) Preferably R¹ is hydroxy, chloro, fluoro, methanesulfonyloxy, amino, azido, methoxy, methylthio, methylaminocarbonyloxy, or of the formula -NHC(=O)R^a wherein R^a is hydrogen, methoxy, amino, chloromethyl, dichloromethyl, cyanomethyl, methoxymethyl,
20 acetylmethyl, methylamino, dimethylamino or (1-4C)alkyl or R¹ is of the formula -NHS(O)_n(1-4C)alkyl wherein n is 0, 1 or 2.

(b) More preferably R¹ is hydroxy, chloro, fluoro, methanesulfonyloxy, or of the formula -NHC(=O)R^a wherein R^a is hydrogen, methoxy, amino, chloromethyl, dichloromethyl, cyanomethyl, methoxymethyl, acetylmethyl or (1-4C)alkyl or R¹ is of the
25 formula -NHS(O)_n(1-4C)alkyl wherein n is 0, 1 or 2.

(c) Yet more preferably R¹ is hydroxy, or of the formula -NHC(=O)R^a wherein R^a is (1-4C)alkyl or R¹ is of the formula -NHS(O)_n(1-4C)alkyl wherein n is 0, 1 or 2.

(d) Yet more preferably R¹ is of the formula -NHC(=O)(1-4C)alkyl or -NHS(O)_n(1-4C)alkyl.

30 (e) Yet more preferably R¹ is of the formula -NHC(=O)(1-4C)alkyl.

(f) Most preferably R¹ is acetamido.


- (g) In another aspect R^1 is hydroxy.
- (h) Preferably, at least one of R^2 and R^3 is hydrogen.
- (i) Preferably R^6 is hydrogen or (1-4C)alkyl.
- (j) Most preferably one of R^2 and R^3 is hydrogen and the other is fluoro.
- 5 (k) Preferably >A-B- is of the formula $>C=CH-$ or $>CHCH_2-$.
- (l) Most preferably >A-B- is of the formula $>C=CH-$.
- (m) Preferably R^4 and R^5 are independently hydrogen, (1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl, hydroxymethyl, (1-4C)alkoxymethyl or carbamoyl.
- (n) More preferably, R^4 and R^5 are independently hydrogen, AR-oxymethyl or
- 10 AR-thiomethyl. Especially preferred is AR when it is optionally substituted phenyl, phenyl(1-4C)alkyl, naphthyl, furan, pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole or thiophene.
- (o) Most preferably R^4 or R^5 is hydrogen.
- (p) Preferably R^6 is hydrogen or (1-4C)alkyl.
- 15 (q) More preferably R^6 is hydrogen or methyl.
- (r) Most preferably R^6 is hydrogen.
- (s) Preferably D is O, S or of the formula NR^7 .
- (t) Preferably D is O.
- (u) Preferred substituents for phenyl and carbon atoms in heteroaryl (mono- and
- 20 bicyclic) ring systems in R^7 and R^{11} include halo, (1-4C)alkyl, hydroxy, nitro, amino, cyano, (1-4C)alkylS(O)_p- and (1-4C)alkoxy.
- (v) Preferred optional substituents for (1-6C)alkyl in R^{11} are hydroxy, cyano, amino, (1-4C)alkylamino, di((1-4C)alkyl)amino, (1-4C)alkylS(O)_p (wherein p is 1 or 2), carboxy, (1-4C)alkoxycarbonyl, (1-4C)alkoxy, piperazino or morpholino.
- 25 (w) Preferred optional substituents for (1-6C)alkyl in R^{12} are hydroxy, (1-4C)alkoxy, cyano, amino, (1-4C)alkylamino, di(C₁₋₂alkyl)amino, (1-4C)alkylS(O)_p (wherein p is 1 or 2).
- (x) Preferably the ring systems in AR are unsubstituted.
- (y) Preferably the 5/6 or 6/6 bicyclic ring system in R^4 , R^5 , R^7 or R^{10} is unsubstituted.
- (z) Preferably 5- or 6-membered heteroaryl rings in R^4 , R^5 , R^7 , R^{10} or R^{11} are
- 30 unsubstituted.
- (a1) Preferably 5- or 6-membered heteroaryl in R^{11} is pyridyl or imidazol-1-yl.

- 15 -

- (b1) Preferably R^{12} is (1-6C)alkyl. Most preferably R^{12} is tert-butyl or methyl.
- (c1) Preferably R^{13} is cyano or fluoro.
- (d1) Preferably R^{14} is hydrogen.
- (e1) Preferably R^{10} is (1-4C)alkoxycarbonyl, hydroxy(1-4C)alkyl, (1-4C)alkyl,
- 5 (1-4C)alkylamino, dimethylamino(1-4C)alkyl, (1-4C)alkoxymethyl, (1-4C)alkanoylmethyl, (1-4C)alkanoyloxy(1-4C)alkyl, (1-5C)alkoxy or 2-cyanoethyl.
- (f1) More preferably R^{10} is methoxycarbonyl, hydroxymethyl, methyl, methylamino, dimethylaminomethyl, methoxymethyl, acetoxymethyl, methoxy, methylthio, naphthyl, tert-butoxy or 2-cyanoethyl.
- 10 (g1) Preferably R^7 is hydrogen, cyano, benzyl, pyrimidyl, imidazolyl, triazolyl or of the formula $R^{10}CO-$ or $R^{10}SO_2-$.
- (h1) Preferably R^7 is hydrogen, cyano, benzyl, methoxycarbonyl, tert-butoxycarbonyl, hydroxyacetyl, dimethylaminoacetyl, acetyloxymethylcarbonyl, methoxyacetyl, methoxalyl, methylcarbamoyl or methanesulfonyl.
- 15 (i1) Preferably when R^{10} is benzopyranone it is 4-oxo-benzopyran-2-yl.
- (j1) Preferably CY is naphthoxy, especially naphth-1-oxy or naphth-2-oxy.

Therefore, especially preferred compounds of the present invention are of the

20 formula (IB):




(IB)

- 25 wherein R^1 is hydroxy or acetamido; R^2 is hydrogen or fluoro and R^4 and R^5 are independently hydrogen, hydroxy, bromo, (1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl, hydroxymethyl, (1-4C)alkoxymethyl or carbamoyl; and pharmaceutically-acceptable salts thereof.

Of the above especially preferred compounds of the formula (IB), particularly preferred compounds are those wherein R¹ is acetamido; R² is hydrogen or fluoro and R⁴ and R⁵ are independently hydrogen, hydroxy, bromo, (1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl, hydroxymethyl, (1-4C)alkoxymethyl or carbamoyl; and
 5 pharmaceutically-acceptable salts thereof.

Further especially preferred compounds of the invention are of the formula (IB) wherein R¹ is acetamido; R² is hydrogen or fluoro and R⁴ and R⁵ are independently hydrogen, AR-oxymethyl or AR-thiomethyl (wherein AR is phenyl, phenyl(1-4C)alkyl, naphthyl, furan, pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole,
 10 isothiazole, thiazole or thiophene); and pharmaceutically-acceptable salts thereof.

Further, especially preferred compounds of the invention are of the formula (IC):



(IC)

15 wherein R¹ is hydroxy or acetamido; R² is hydrogen or fluoro; R⁴ and R⁵ are independently hydrogen. AR-oxymethyl or AR-thiomethyl (wherein AR is phenyl, phenyl(1-4C)alkyl, naphthyl, furan, pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole or thiophene), hydroxy, bromo, (1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl, hydroxymethyl, (1-4C)alkoxymethyl or carbamoyl and R⁷ is cyano,
 20 pyrimidin-2-yl, tetrazol-5-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl or R⁷ is of the formula R¹⁰CO- or R¹⁰SO₂- (wherein R¹⁰ is hydrogen, (1-4C)alkyl [optionally substituted by hydroxy, (1-4C)alkoxy, (1-4C)alkylS(O)_q, (1-4C)alkylamino, (1-4C)alkanoyl, naphthoxy, (2-6C)alkanoylamino or (1-4C)alkylS(O)_pNH- wherein p is 1 or 2 and q is 0, 1 or 2], phenyl, naphthyl, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole,
 25 isothiazole, thiazole, thiophene, benzofuran, benzoimidazole, benzothiophene, benzisothiazole, benzoxazole, benzisoxazole, pyridoimidazole, pyrimidoimidazole, quinoline, quinoxaline, quinazoline, phthalazine, cinnoline or naphthyridine, or R¹⁰ is of the formula

$R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is $(1-6C)alkyl$, or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is $(1-6C)alkoxy$; and pharmaceutically-acceptable salts thereof.

Of the above especially preferred compounds of the invention of the formula (IC), particularly preferred compounds are those wherein R^1 is acetamido; R^2 is hydrogen or fluoro;
 5 R^4 and R^5 are independently hydrogen, AR-oxyethyl or AR-thioethyl [wherein AR is phenyl, phenyl(1-4C)alkyl, naphthyl, furan, pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole or thiophene], hydroxymethyl, (1-4C)alkoxyethyl or carbamoyl and R^7 is cyano, pyrimidin-2-yl, tetrazol-5-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl, or R^7 is of the formula $R^{10}CO-$ or $R^{10}SO_2-$
 10 (wherein R^{10} is hydrogen, (1-4C)alkyl [optionally substituted by hydroxy, (1-4C)alkylS(O)_q, (1-4C)alkanoyl or naphthoxy wherein q is 0, 1 or 2], phenyl, naphthyl, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole, thiophene, benzofuran, benzoimidazole, benzothiophene, benzisothiazole, benzoxazole, benzisoxazole, pyridoimidazole, pyrimidoimidazole, quinoline, quinoxaline, quinazoline, phthalazine, cinnoline or naphthyridine, or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is $(1-6C)alkyl$, or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is $(1-6C)alkoxy$; and pharmaceutically-acceptable salts thereof.

Of the above particularly preferred compounds of the invention of the formula (IC), especially preferred compounds are those wherein R^1 is acetamido; R^2 is hydrogen or fluoro;
 20 R^4 and R^5 are hydrogen and R^7 is cyano, pyrimidin-2-yl, tetrazol-5-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl, or R^7 is of the formula $R^{10}CO-$ or $R^{10}SO_2-$ (wherein R^{10} is hydrogen, (1-4C)alkyl [optionally substituted by hydroxy, (1-4C)alkylS(O)_q or (1-4C)alkanoyl wherein q is 0, 1 or 2], phenyl, naphthyl, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole, thiophene, benzofuran,
 25 benzoimidazole, benzothiophene, benzisothiazole, benzoxazole, benzisoxazole, pyridoimidazole, pyrimidoimidazole, quinoline, quinoxaline, quinazoline, phthalazine, cinnoline or naphthyridine, or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is $(1-6C)alkyl$, or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is $(1-6C)alkoxy$; and pharmaceutically-acceptable salts thereof.

30 Of the above particularly preferred compounds of the invention of the formula (IC), further especially preferred compounds are those wherein R^1 is acetamido; R^2 is hydrogen or

fluoro; R^4 and R^5 are independently hydrogen, AR-oxymethyl or AR-thiomethyl (wherein AR is phenyl, phenyl(1-4C)alkyl, naphthyl, furan, pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole or thiophene), hydroxymethyl, (1-4C)alkoxymethyl or carbamoyl and R^7 is cyano, pyrimidin-2-yl, tetrazol-5-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl, or R^7 is of the formula $R^{10}CO-$ (wherein R^{10} is hydrogen, (1-4C)alkyl [optionally substituted by hydroxy, (1-4C)alkylS(O)_q or (1-4C)alkanoyl wherein q is 0, 1 or 2], phenyl, naphthyl, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole, thiophene, benzofuran, benzoimidazole, benzothiophene, benzisothiazole, benzoxazole, benzisoxazole, pyridoimidazole, pyrimidoimidazole, quinoline, quinoxaline, quinazoline, phthalazine, cinnoline or naphthyridine, or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is (1-6C)alkyl), or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is (1-6C)alkoxy; and pharmaceutically-acceptable salts thereof.

Of the above particularly preferred compounds of the invention of the formula (IC), chiefly preferred compounds are those wherein R^1 is acetamido; R^2 is hydrogen or fluoro; R^4 and R^5 are independently hydrogen, AR-oxymethyl or AR-thiomethyl (wherein AR is phenyl, phenyl(1-4C)alkyl, naphthyl, furan, pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole or thiophene), hydroxymethyl, (1-4C)alkoxymethyl or carbamoyl and R^7 is cyano, pyrimidin-2-yl, tetrazol-5-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl, or R^7 is of the formula $R^{10}SO_2-$ (wherein R^{10} is hydrogen, (1-4C)alkyl [optionally substituted by hydroxy, (1-4C)alkylS(O)_q or (1-4C)alkanoyl wherein q is 0, 1 or 2], phenyl, naphthyl, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole, thiophene, benzofuran, benzoimidazole, benzothiophene, benzisothiazole, benzoxazole, benzisoxazole, pyridoimidazole, pyrimidoimidazole, quinoline, quinoxaline, quinazoline, phthalazine, cinnoline or naphthyridine, or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is (1-6C)alkyl), or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is (1-6C)alkoxy; and pharmaceutically-acceptable salts thereof.

Of the above further especially preferred compounds of the invention of the formula (IC), particular especially preferred compounds are those wherein R^1 is acetamido; R^2 is hydrogen or fluoro; R^4 and R^5 are hydrogen; and R^7 is pyrimidin-2-yl, 2-cyanoethenyl,

2-cyano-2-((1-4C)alkyl)ethenyl, or R^7 is of the formula $R^{10}CO-$ (wherein R^{10} is hydrogen or (1-4C)alkyl [optionally substituted by hydroxy or (1-4C)alkylS(O)_q wherein q is 0, 1 or 2], or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is (1-6C)alkyl), or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is (1-6C)alkoxy; and pharmaceutically-acceptable salts thereof.

5 Of the above chiefly preferred compounds of the invention of the formula (IC), particular chiefly preferred compounds are those wherein R^1 is acetamido; R^2 is hydrogen or fluoro; R^4 and R^5 are hydrogen; and R^7 is pyrimidin-2-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl, or R^7 is of the formula $R^{10}SO_2-$ (wherein R^{10} is hydrogen or (1-4C)alkyl [optionally substituted by hydroxy or (1-4C)alkylS(O)_q wherein q is 0, 1 or 2], or R^{10} is of the
10 formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is (1-6C)alkyl), or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is (1-6C)alkoxy; and pharmaceutically-acceptable salts thereof.

Of the above particular especially preferred compounds of the invention of the formula (IC), the most preferred are those wherein R^1 is acetamido; R^2 is hydrogen or fluoro; R^4 and R^5 are hydrogen; and R^7 is pyrimidin-2-yl, or R^7 is of the formula $R^{10}CO-$ (wherein R^{10}
15 is hydrogen or (1-4C)alkyl [optionally substituted by hydroxy or (1-4C)alkylS(O)_q wherein q is 0, 1 or 2], or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is (1-6C)alkyl), or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is (1-6C)alkoxy; and pharmaceutically-acceptable salts thereof.

Of the above particular chiefly preferred compounds of the invention of the formula
20 (IC), the most preferred are those wherein R^1 is acetamido; R^2 is hydrogen or fluoro; R^4 and R^5 are hydrogen; and R^7 is pyrimidin-2-yl, or R^7 is of the formula $R^{10}SO_2-$ (wherein R^{10} is hydrogen or (1-4C)alkyl [optionally substituted by hydroxy or (1-4C)alkylS(O)_q wherein q is 0, 1 or 2], or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is (1-6C)alkyl), or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is (1-6C)alkoxy; and pharmaceutically-acceptable
25 salts thereof.

Particular compounds of the present invention are:

N-((5S)-3-(4-(1-tert-butoxycarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(4-(1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

30 N-((5S)-3-(4-(1-methoxycarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide:.

- N-((5S)-3-(4-(1-methylsulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(4-(1-hydroxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 5 N-((5S)-3-(4-(1-dimethylaminoacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(4-(2,3-dihydropyran-4-yl)-phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(4-(2,3-dihydrothiapyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(4-(2,3-dihydrooxothiapyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 10 N-((5S)-3-(4-(2,3-dihydrodioxothiapyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1-benzyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxo-oxazolidin-5-ylmethyl)acetamide;
- 15 N-((5S)-3-(3-fluoro-4-(1-tertbutoxycarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxo-5-oxazolidinylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1-methoxycarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 20 N-((5S)-3-(3-fluoro-4-(1-methylsulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1-hydroxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1-dimethylaminoacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 25 N-((5S)-3-(3-fluoro-4-(2,3-dihydropyran-4-yl)-phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(2,3-dihydrothiapyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 30 N-((5S)-3-(3-fluoro-4-(2,3-dihydrooxothiapyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

- N-((5S)-3-(4-(2,3-dihydrodioxothiapyrany-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(4-(1-(pyrimidin-2-yl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 5 N-((5S)-3-(4-(1-cyano-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1-(acetoxycetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 5R-hydroxymethyl-3-(3-fluoro-4-(1-benzyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)oxazolidin-2-
- 10 one;
- N-((5S)-3-(3-fluoro-4-(1-(pyrimidin-2-yl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1-cyano-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 15 N-((5S)-3-(3-fluoro-4-(1-methoxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1-methoxalyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1-(N-methylcarbamoyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-
- 20 oxooxazolidin-5-ylmethyl)acetamide;
- and pharmaceutically-acceptable salts thereof.

Other particular compounds of the present invention are

- N-((5S)-3-(4-(1-(2-cyano-2-methylethenyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 25 N-((5S)-3-(4-(1-(hydroxymethylsulfonyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(4-(1-(methylsulfonylaminomethylcarbonyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxo-oxazolidin-5-ylmethyl)acetamide;
- and pharmaceutically-acceptable salts thereof.

- 30 Particularly preferred compounds of the present invention are:

- N-((5S)-3-(4-(1-methylsulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxo-oxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(4-(1-acetyloxymethylcarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 5 N-((5S)-3-(4-(1-hydroxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1-acetyloxymethylcarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1-hydroxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 10 N-((5S)-3-(4-(2,3-dihydro-6H-pyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(2,3-dihydro-6H-pyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(3-fluoro-4-(1-{pyrimid-2-yl}-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 15 N-((5S)-3-(3-fluoro-4-(1-methylsulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(4-(1-{pyrimid-2-yl}-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 20 N-((5S)-3-(3-fluoro-4-(1-acetoacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(4-(1-(naphth-2-ylsulfonyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(4-(1-(naphth-2-oxyacetyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 25 N-((5S)-3-(4-(1-(methylthioacetyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- and pharmaceutically-acceptable salts thereof.

Especially preferred compounds of the invention are:

- 30 N-((5S)-3-(4-(1-methylsulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxo-oxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(4-(1-acetyloxymethylcarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(4-(1-hydroxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

5 N-((5S)-3-(3-fluoro-4-(1-acetyloxymethylcarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(3-fluoro-4-(1-hydroxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(4-(2,3-dihydro-6H-pyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

10 N-((5S)-3-(3-fluoro-4-(2,3-dihydro-6H-pyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

and pharmaceutically-acceptable salts thereof.

Other especially preferred compounds of the invention are:

N-((5S)-3-(3-fluoro-4-(1-{pyrimid-2-yl}-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-

15 oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(3-fluoro-4-(1-methylsulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(4-(1-{pyrimid-2-yl}-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

20 N-((5S)-3-(3-fluoro-4-(1-acetoacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(4-(1-(naphth-2-ylsulfonyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;


N-((5S)-3-(4-(1-(naphth-2-oxyacetyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-25 5-ylmethyl)acetamide;

N-((5S)-3-(4-(1-(methylthioacetyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

and pharmaceutically-acceptable salts thereof.

In a further aspect the present invention provides a process for preparing a

30 compound of formula (I) or a pharmaceutically-acceptable salt thereof. The compounds of formula (I) may be prepared by deprotecting a compound of formula (II):



(II)

5

wherein R^2 , R^3 and R^6 are as hereinabove defined, R^{18} is R^4 or protected R^4 , R^{19} is R^5 or protected R^5 , R^{20} is R^1 or protected R^1 , $>A1-B1-$ is $>A-B-$ or protected $>A-B-$ and D1 is D in which functional groups are optionally protected; and thereafter, if necessary, forming a pharmaceutically-acceptable salt.

10

Protecting groups may be removed by any convenient method as described in the literature or known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so as to effect removal of the protecting group with minimum disturbance of groups elsewhere in the molecule.

Specific examples of protecting groups are given below for the sake of convenience, in which "lower" signifies that the group to which it is applied preferably has 1-4 carbon atoms. It will be understood that these examples are not exhaustive. Where specific examples of methods for the removal of protecting groups are given below these are similarly not exhaustive. The use of protecting groups and methods of deprotection not specifically mentioned is of course within the scope of the invention.

20

A carboxy protecting group may be the residue of an ester-forming aliphatic or araliphatic alcohol or of an ester-forming silanol (the said alcohol or silanol preferably containing 1-20 carbon atoms).

Examples of carboxy protecting groups include straight or branched chain (1-12C)alkyl groups (eg isopropyl, t-butyl); lower alkoxy lower alkyl groups (eg methoxymethyl, ethoxymethyl, isobutoxymethyl; lower aliphatic acyloxy lower alkyl groups, (eg acetoxymethyl, propionyloxymethyl, butyryloxymethyl, pivaloyloxymethyl); lower alkoxycarbonyloxy lower alkyl groups (eg 1-methoxycarbonyloxyethyl,

25

1-ethoxycarbonyloxyethyl); aryl lower alkyl groups (eg p-methoxybenzyl, o-nitrobenzyl, p-nitrobenzyl, benzhydryl and phthalidyl); tri(lower alkyl)silyl groups (eg trimethylsilyl and t-butyldimethylsilyl); tri(lower alkyl)silyl lower alkyl groups (eg trimethylsilylethyl); and (2-6C)alkenyl groups (eg allyl and vinyllethyl).

5 Methods particularly appropriate for the removal of carboxyl protecting groups include for example acid-, metal- or enzymically-catalysed hydrolysis.

 Examples of hydroxy protecting groups include lower alkenyl groups (eg allyl); lower alkanoyl groups (eg acetyl); lower alkoxycarbonyl groups (eg t-butoxycarbonyl); lower alkenyloxycarbonyl groups (eg allyloxycarbonyl); aryl lower alkoxycarbonyl groups (eg
10 benzoyloxycarbonyl, p-methoxybenzyloxycarbonyl, o-nitrobenzyloxycarbonyl, p-nitrobenzyloxycarbonyl); tri lower alkyl/arylsilyl groups (eg trimethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl); aryl lower alkyl groups (eg benzyl) groups; and triaryl lower alkyl groups (eg triphenylmethyl).

 Examples of amino protecting groups include formyl, aralkyl groups (eg benzyl and
15 substituted benzyl, eg p-methoxybenzyl, nitrobenzyl and 2,4-dimethoxybenzyl, and triphenylmethyl); di-p-anisylmethyl and furylmethyl groups; lower alkoxycarbonyl (eg t-butoxycarbonyl); lower alkenyloxycarbonyl (eg allyloxycarbonyl); aryl lower alkoxycarbonyl groups (eg benzyloxycarbonyl, p-methoxybenzyloxycarbonyl, o-nitrobenzyloxycarbonyl, p-nitrobenzyloxycarbonyl); trialkylsilyl (eg trimethylsilyl and
20 t-butyldimethylsilyl); alkylidene (eg methylidene); benzylidene and substituted benzylidene groups.

 Methods appropriate for removal of hydroxy and amino protecting groups include, for example, acid-, metal- or enzymically-catalysed hydrolysis, for groups such as o-nitrobenzyloxycarbonyl, photolytically and for groups such as silyl groups, fluoride.

25 Examples of protecting groups for amide groups include aralkoxymethyl (eg. benzyloxymethyl and substituted benzyloxymethyl); alkoxymethyl (eg. methoxymethyl and trimethylsilylethoxymethyl); tri alkyl/arylsilyl (eg. trimethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl); tri alkyl/arylsilyloxymethyl (eg. t-butyldimethylsilyloxymethyl, t-butyldiphenylsilyloxymethyl); 4-alkoxyphenyl (eg. 4-methoxyphenyl); 2,4-di(alkoxy)phenyl
30 (eg. 2,4-dimethoxyphenyl); 4-alkoxybenzyl (eg. 4-methoxybenzyl); 2,4-di(alkoxy)benzyl


(eg. 2,4-di(methoxy)benzyl); and alk-1-enyl (eg. allyl, but-1-enyl and substituted vinyl eg. 2-phenylvinyl).

Aralkoxymethyl, groups may be introduced onto the amide group by reacting the latter group with the appropriate aralkoxymethyl chloride, and removed by catalytic hydrogenation. Alkoxymethyl, tri alkyl/arylsilyl and tri alkyl/silyl groups may be introduced by reacting the amide with the appropriate chloride and removing with acid, or in the case of the silyl containing groups fluoride ions. The alkoxyphenyl and alkoxybenzyl groups are conveniently introduced by arylation or alkylation with an appropriate halide and removed by oxidation with ceric ammonium nitrate. Finally alk-1-enyl groups may be introduced by reacting the amide with the appropriate aldehyde and removed with acid.

For further examples of protecting groups see one of the many general texts on the subject, for example, 'Protective Groups in Organic Synthesis' by Theodora Green (publisher: John Wiley & Sons).


In another aspect of the present invention the compounds of the formulae (I) and (II) and pharmaceutically-acceptable salts thereof can be prepared:

- (a) by modifying a substituent in or introducing a substituent into another compound of formula (I) or (II);
- (b) when R^1 or R^{20} is of the formula $-NHS(O)_n(1-4C)alkyl$, wherein n is 1 or 2, by oxidising a compound of the formula (I) or (II) wherein n is 0 or, when n is 2 by oxidising a compound of the formula (I) or (II) wherein n is 1;
- (c) when R^1 or R^{20} is of the formula $-NHC(=O)R^b$ or $NHS(O)_n(1-4C)alkyl$, introducing the group $-C(=O)R^b$ or $-S(O)_n(1-4C)alkyl$ into a compound of the formula (III);




(III)

- (d) when R^1 or R^{20} is hydroxy, by reacting a compound of the formula (V) with a compound of formula (VI):




(V)




(VI)

5

(e) when $>A1-B1-$ is $>C=CR^{a1}-$, by reacting a compound of the formula (VII) with a compound of the formula (VIII):



(VII)




(VIII)

10 (f) when $>A1-B1-$ is $>CHCH(R^{a1})-$, by catalytic hydrogenation of a compound of the formula (I) or (II) wherein $>A1-B1-$ is $>C=CR^{a1}-$;


(g) when $>A1-B1-$ is $>C=CR^{a1}-$, by elimination of the elements of water, or $HOCOR^{23}$, or $HOSO_2R^{24}$ from a compound of the formula (IX) (ie. when R^{25} is $-H$, $-COR^{23}$ or $-SO_2R^{24}$).

15




(IX)

(h) when D is NR^7 and R^7 is $\text{R}^{10}\text{CO}-$ or $\text{R}^{10}\text{S(O)}_n-$, wherein n is 2, by reaction of a compound of formula (X) with a compound of the formula (XI) or (XII), wherein n is 2:



(i) when R^1 or R^{20} is azido, by reacting a compound of the formula (XIII) with a source of azide:



(j) when R^1 or R^{20} is amino, by reducing a compound of the formula (I) or (II) wherein
15 R^1 or R^{20} is azido;

(k) when R^1 or R^{20} is chloro, fluoro, (1-4C)alkanesulfonyloxy, or (1-4C)alkylaminocarbonyloxy, or R^{20} is of the formula $-\text{N}(\text{CO}_2\text{R}^{27})\text{CO}(1-4\text{C})\text{alkyl}$; from a compound of the formula (I) and (II) wherein R^1 or R^{20} is hydroxy; or

(l) when R^1 or R^{20} is chloro, (1-4C)alkylthio or (1-4C)alkoxy, from a compound of the
20 formula (XIII);

wherein R^2 , R^3 , R^6 and R^{20} are as hereinabove defined; R^{11} is R^a or protected R^a ; R^{21} is (1-6C)alkyl or benzyl; R^{22} is of the formula (1-4C)alkyl or $-\text{S(O)}_n(1-4\text{C})\text{alkyl}$; R^{23} is (1-4C)alkyl; R^{24} is an optionally substituted phenyl group; R^{25} is hydrogen, $-\text{COR}^{23}$

or $-\text{SO}_2$, R^{24} ; R^{26} is mesyloxy or tosyloxy; R^{27} is (1-4C)alkyl or benzyl; n is 0, 1 or 2 unless otherwise stated above; L^1 is an iodo or triflate leaving group; L^2 is a leaving group, such as, for example, hydroxy or chloro; and Y is a trialkyltin residue or a boronate acid or ester residue;

- 5 and thereafter if necessary: i) removing any protecting groups;
ii) forming a pharmaceutically-acceptable salt.

Methods for converting substituents into other substituents are known in the art. For example an alkylthio group may be oxidised to an alkylsulfinyl or alkylsulfonyl group, a cyano group reduced to an amino group, a nitro group reduced to an amino group, a hydroxy group
10 alkylated to a methoxy group, a hydroxy group thiomethylated to an arylthiomethyl or a heteroarylthiomethyl group (see, for example, Tet.Lett., 585, 1972), a carbonyl group converted to a thiocarbonyl group (eg. using Lawesson's reagent) or a bromo group converted to an alkylthio group.

Compounds of the formula (I) or (II) wherein R^1 or R^{20} is $-\text{NHS}(\text{O})_n$ (1-4C)alkyl
15 can be prepared by oxidising a compound of the formula (I) or (II) with standard reagents known in the art for the oxidation of a thio group to a sulfinyl or sulfonyl group. For example, a thio group may be oxidised to a sulfinyl group with a peracid such as m-chloroperoxybenzoic acid and oxidising agents such as potassium permanganate can be used to convert a thio group to a sulfonyl group. Compounds of the formula (I) or (II)
20 wherein R^1 or R^{20} is $-\text{NHS}(1-4\text{C})\text{alkyl}$ can be prepared by reacting compounds of the formula (III) with a reagent such as (1-4C)alkylSCl.

When R^b is (1-4C)alkyl, the group $-\text{C}(=\text{O})(1-4\text{C})\text{alkyl}$ may be introduced into a compound of the formula (III) by standard acetylation procedures. For example, the amino group may be acetylated to give an acetamido group using the Schotten-Baumann procedure
25 i.e. reacting the compound of the formula (I) or (II) wherein R^1 or R^{20} is amino with acetic anhydride in aqueous sodium hydroxide and THF in a temperature range of 0°C to ambient temperature. Preferably the acylation is carried out in situ following the catalytic hydrogenation of a compound of the formula (I) or (II) wherein R^1 or R^{20} is azido, by performing the hydrogenation in the presence of acetic anhydride.

30 When R^b is hydrogen, the $-\text{CHO}$ group may be introduced into the compound of the formula (III) by reacting the latter compound with formic acetic anhydride, in an inert organic

solvent such as THF, in a temperature range of 0°C to ambient temperature, or by reacting it with ethyl formate in an inert organic solvent in the temperature range of 50-100°C.

When R^b is (1-4C)alkoxy, the -COO(1-4C)alkyl group may be introduced into the compound of the formula (III) by reacting the latter compound with (1-4C)alkyl
5 chloroformate, in the presence of an organic base such as triethylamine, in an organic solvent such as dichloromethane and in a temperature range of 0°C to ambient temperature.

When R^b is amino, the -CONH₂ group may be introduced into the compound of the formula (III) by reacting the latter compound either with potassium cyanate in aqueous acid (eg hydrochloric acid) in a temperature range of ambient temperature to 40°C or with phenyl
10 carbamate in glyme at reflux.

When R^b is chloromethyl, dichloromethyl, cyanomethyl or methoxymethyl, the -C(=O)R^b group may be introduced into the compound of the formula (III) by reacting the latter compound with the appropriate acid chloride under standard conditions. The acid
chloride may be prepared from the appropriate acid. When R^b is acetylmethyl, the -C(=O)R^b
15 group may be introduced into the amino compound by reacting the latter compound with diketene, in an inert organic solvent such as THF, in a temperature range of 0°C to ambient temperature.

Alternatively, the compound of the formula (III) may be reacted with the appropriate acid anhydride, in dichloromethane or THF, in the presence of an organic base such as
20 triethylamine and in a temperature range of 0°C to ambient temperature, or the amino compound may be reacted with the appropriate acid in the presence of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and an organic base such as triethylamine, in an organic solvent such as dichloromethane, in a temperature range of 0°C to ambient temperature.

25 When R^b is methylamino, the -CONHMe group may be introduced into the compound of the formula (III) by reacting the latter compound with methyl isocyanate in an organic solvent such as THF or acetonitrile, in a temperature range of 0°C to ambient temperature.


When R^b is dimethylamino, the -CONMe₂ group may be introduced into the
30 compound of the formula (III) by reacting the latter compound with dimethylcarbonyl

chloride and triethylamine in an organic solvent such as THF or acetonitrile, in a temperature range of 0°C to ambient temperature.

Standard reaction conditions for the conversion of a compound of the formula (III) to a compound of the formula (I) or (II) wherein R¹ or R²⁰ is sulfonamido are known in the art.

5 For example, a compound of the formula (III) could be converted to a compound of the formula (I) or (II) wherein R¹ or R²⁰ is (1-4C)alkylSO₂NH- by reacting the former compound with a sulfonyl chloride, for example, mesyl chloride, in a mild base such as pyridine or triethylamine.


Alternatively compounds of the formula (I) or (II) wherein R¹ or R²⁰ is
10 (1-4C)alkylSO₂NH- or (1-4C)alkylSONH- may be prepared by reacting a compound of the formula (III) with a compound of the formula (IV):



(IV)

15

The compound of the formula (IV) may be prepared by oxidising a compound of the formula (IVA):



(IVA)


20 with standard oxidising agents known for the conversion of a thio group to a sulfinyl or sulfonyl group.

Compounds of the formula (IVA) can be prepared by reacting phthalimide with an alkylthiochloride ((1-4C)alkylSCl).

A compound of the formula (III) may be prepared as described in process (j).

Compounds of the formulae (V) and (VI) are conveniently reacted together in the presence of a strong base such as butyl lithium, lithium hexamethyldisilazide, sodium hydride, or lithium diisopropylamide. The reaction is conveniently carried out in an inert solvent such as tetrahydrofuran (THF), dimethylformamide (DMF), N,N'-dimethylpropyleneurea (DMPU) or N-methylpyrrolidone in a temperature range of -78°C to -50°C for the deprotonation and cyclisation. Suitable values for R²¹ include ethyl, butyl and benzyl and suitable values for R²² include ethyl and n-propyl, preferably n-propyl.


10 A compound of the formula (V) is conveniently prepared by reacting a chloroformate of the formula (ClCOOR²¹) with a compound of the formula (VA):



15 (VA)

wherein R² - R⁵ and >A1-B1- are as hereinabove defined. The reaction is conveniently carried out in the presence of an inorganic or organic base such as sodium bicarbonate or an amine base such as dimethylaniline, the former in a solvent such as acetone/water and the latter in an organic solvent such as THF, toluene, DMF or acetonitrile.

20 A compound of the formula (VA) wherein >A1-B1- is >C=CH-, may be prepared by reacting a compound of the formula (VIII) with a compound of the formula (VB).



The reaction between compounds of the formulae (VIII) and (VB) wherein L^1 is bromo or iodo may be carried out by treating (VB) with an organolithium species such as 1-butyl methyl in an inert solvent such as THF at a low temperature, such as -78°C , followed by the addition of an anhydrous zinc halide such as zinc chloride, in a temperature range of 0°C to ambient temperature, to generate the organozinc chloride (VB), wherein L^1 is ZnCl . Treatment of the organozinc chloride in situ with a compound of the formula (VIII) followed by a suitable palladium (0) catalyst such as $\text{Pd}(\text{PPh}_3)_4$, in the temperature range of 0°C to ambient temperature, results in the cross-coupled product (VA) after brief treatment with dilute acid to hydrolyse the 'stabase' protected amine.


- 10 A compound of the formula (VB) may be prepared by treatment of p-iodo or p-bromoaniline with the 'stabase' reagent (1,2-bis(chlorodimethylsilyl)ethane) in the presence of an organic base such as triethylamine.

The reaction between compounds of the formulae (VII) and (VIII), wherein Y is trialkyltin and L^1 is triflate is conveniently carried out in the presence of a palladium (0) catalyst such as $\text{Pd}(\text{PPh}_3)_4$ or $\text{Pd}(\text{dba})_3$, in a temperature range of $0-115^\circ\text{C}$. Preferably the trialkyltin group is trimethyltin. A suitable value for L^1 is iodo or trifluoromethylsulfonyloxy.

When Y is a boronate acid or ester, the reaction may be carried out under conditions known for the Suzuki reaction i.e. in the presence of a palladium (0) catalyst such as $\text{Pd}(\text{PPh}_3)_4$ or $\text{Pd}(\text{dba})_3$, in a water-miscible organic solvent such as dimethylformamide or 1,2-dimethoxyethane and in the presence of a mild base such as sodium acetate or sodium bicarbonate which is added in water. The reaction is then heated to 80°C . Alternatively, silver oxide may be used in place of the base, in which case the reaction may be carried out at a lower temperature. When Y is a boronate ester, preferably L^1 is iodo. Suitable boronate esters include lower alkyl and cyclic boronate esters.

- 25 A compound of the formula (VII) wherein Y is trimethylstannyl may be prepared by methods known in the art (for example by using methods similar to those described in Patent Application No. WO 9413649 from a compound of the formula (VII) wherein Y is iodo or bromo). Alternatively compounds of the formula (VII) wherein Y is a cyclic boronate ester as in (VIIA):


- 34 -



(VIIA)

may be prepared from a compound of the formula (VII) wherein Y is iodo or bromo, by
 5 sequential treatment with a suitable Pd catalyst such as PdCl₂(dppf), potassium acetate and the
 pinacol ester of diboron in a polar solvent such as DMSO (for example see J.Org.Chem.,
 1995, 60, 7508-7510).

A compound of the formula (VII), wherein Y is iodo may be prepared by reacting a
 compound of the formula (VIIB) with iodine monochloride in the presence of trifluoroacetic
 10 acid or with iodine and silver triflate:




(VIIB)

When Y is bromo, a compound of the formula (VII) may be prepared by brominating
 15 a compound of the formula (VIIB) using standard bromination methods. For example, by
 reacting a compound of the formula (VIIB) with N-bromosuccinimide or bromine.

A compound of the formula (VIIB) may be prepared by forming the oxazolidinone
 ring from the amino group of a compound of the formula (VIIC) using a similar method to
 that described for the preparation of a compound of the formula (I) or (II) from a compound of
 20 the formula (VA):


- 35 -



(VIIC)

5 The resulting compound of the formula (VIIB) in which R^{20} is hydroxy may be converted to other compounds of the formula (VIIB) using similar methods to those described for the formation of a compound of the formula (I) or (II) from a compound of the formula (I) or (II) wherein R^1 or R^{20} is hydroxy, via a compound of the formula (III).

10 A compound of the formula (VIII) wherein $D1$ is $R^{10}CON-$, S or O and L^1 is triflate may be prepared by treating a compound of the formula (VIIIA) with lithium diisopropylamide in an inert solvent such as THF, at a low temperature, for example -78°C , followed by N-phenyl triflamide (for example, see methods described in Synthesis, 993-95 (1991)).



(VIIIA)

15 Alternatively, a compound of the formula (VIII) wherein L^1 is iodo may be prepared by treating a hydrazone of a compound of formula (VIIIA) with iodine in the presence of
20 triethylamine (for example see methods detailed in Tet. Letts., 24, 1605-1608 (1983)).

Compounds of the formula (VIII) wherein $D1$ is $R^{14}CH(R^{13})(CH_2)_m-N<$, aryl- $N<$ or heteroaryl(mono or bicyclic)- N can be prepared by elaboration of the piperidone ring from the appropriate aryl- or heteroarylamine, by reaction with ethyl acrylate to give the

corresponding diethylarylimino-bb'-dipropionate, which can then be cyclised under Diekmann conditions to give corresponding piperidone β -ketoester, followed by decarboxylation with heating in acid (see methods described in J.Chem.Soc., 5110-5118 (1962)).

5 Alternatively, a compound of the formula (VIII) wherein D1 is heteroaryl-N< may be prepared by reacting an appropriately substituted heterocycle containing a leaving group such as chloro, bromo or iodo with the appropriate 4-piperidone at an elevated temperature, in an inert solvent and optionally with an acid trapping agent.

Suitable catalysts for the catalytic hydrogenation of a compound of the formula (I) or
10 (II) wherein $>A1-B1-$ is $>C=C(R^{a1})-$ include Raney nickel, platinum metal or its oxide, rhodium, zinc oxide, palladium-on-charcoal and Wilkinson's catalyst $(RhCl(Ph_3P)_3)$. Catalytic hydrogenation is conveniently carried out in the temperature range $0^\circ C$ to $150^\circ C$, but preferably at ambient temperature and pressure, unless Wilkinson's catalytic is used in which case a temperature of approximately $50^\circ C$ and pressure of approximately 50 atmospheres are
15 preferable.

A compound of the formula (IX) may be prepared by reacting an intermediate of the formula (VB) with magnesium to form a Grignard reagent, or alternatively with n-butyl lithium to form a lithiated species (as above), and then reacting the Grignard reagent or lithiated species with a compound of formula (VIII A). The product (VA), wherein $>A1-B1-$
20 is of the formula $>C(OH)CH(R^a)-$ may then be elaborated as previously detailed for the compound of the formula (V), but with optional protection of the hydroxyl group.

The dehydration of a compound of the formula (IX) to give a compound of formula (I) or (II) wherein $>A1-B1-$ is of the formula $>C=CR^{a1}-$ may be carried out using agents such as polyphosphoric acid, trifluoroacetic acid, trifluoroacetic anhydride, p-toluenesulfonic
25 acid, sulfuric acid, thionyl chloride etc., in an inert solvent such as toluene, and at elevated temperatures. Suitable protection of the group R^{20} may be necessary as appropriate.

A compound of the formula (I) or (II) wherein $>A1-B1-$ is of the formula $>CHCH(R^{a1})-$ may be prepared from a compound of the formula (I) or (II) wherein $>A1-B1-$ is $>C=CR^{a1}-$, by catalytic hydrogenation, using a suitable catalyst such as palladium-on-
30 carbon in an appropriate inert or acidic solvent such as acetic acid. Where an optically active form of compounds of the formula (VI) is used in previous steps, reduction of the $>A1-B1-$

double bond will produce diastereoisomers which may be separated. Where a particular diastereoisomer is of choice, a chiral asymmetry-inducing catalyst for the reduction can be used.

The reaction between a compound of the formula (X) and (XI) or (XII) is
5 conveniently carried out under similar conditions to those described for the acetylation or sulfonylation of a compound of the formula (III).

A compound of the formula (I) or (II) wherein R¹ or R²⁰ is azido may be prepared, for example, by reacting a compound of the formula (XIII) with sodium azide in an inert solvent such as DMF in a temperature range of ambient to 100°C, normally in the region of
10 75°C - 85°C. A compound of the formula (XIII) may be prepared by converting the hydroxy group in a compound of the formula (I) or (II) wherein R¹ or R²⁰ is hydroxy into a tosyloxy or mesyloxy group by standard methods known in the art. For example, by reacting the compound of the formula (I) or (II) with tosyl chloride or mesyl chloride in the presence of a mild base such as triethylamine, or pyridine.

15 Suitable reducing agents for reducing azido to amino in a compound of the formula (I) or (II) include triethylamine/hydrogen sulfide, triphenylphosphine or phosphite ester, or hydrogen in the presence of a catalyst. More specifically the reduction of the azido group may be carried out by heating it in an aprotic solvent, such as 1,2-dimethoxyethane, in the presence of P(OMe)₃, and subsequently heating in 6N aqueous hydrochloric acid, or reacting it
20 with hydrogen in the presence of palladium on carbon in a solvent such as DMF or ethyl acetate. For further details on the reduction of azides to amines see USP 4,705,799. The azido compound may be reduced and converted to a compound of the formula (I) or (II), wherein R¹ or R¹⁰ is acetamido, in situ using acetic anhydride in DMF.

A compound of the formula (I) or (II) wherein R¹ or R²⁰ is fluoro may be prepared
25 by reacting a compound of the formula (I) or (II) wherein R¹ or R²⁰ is hydroxy (hydroxy compound) with a fluorinating agent such as diethylaminosulfur trifluoride in an organic solvent such as dichloromethane in the temperature range of 0°C to ambient temperature.

When R¹ or R²⁰ is chloro, the compound of the formula (I) or (II) may be formed by reacting the hydroxy compound with a chlorinating agent. For example, by reacting the
30 hydroxy compound with thionyl chloride, in a temperature range of ambient temperature to reflux, optionally in a chlorinated solvent such as dichloromethane or by reacting the hydroxy

compound with carbon tetrachloride/triphenyl phosphine in dichloromethane, in a temperature range of 0°C to ambient temperature.

The (1-4C)alkanesulfonyloxy compound may be prepared by reacting the hydroxy compound with (1-4C)alkanesulfonyl chloride in the presence of a mild base such as
5 triethylamine or pyridine.

The (1-4C)alkylaminocarbonyloxy compound may be prepared by reacting the hydroxy compound with (1-4C)alkyl cyanate in an organic solvent such as THF or acetonitrile, in the presence of triethylamine, in a temperature range of 0°C to 50°C.

A compound of the formula (II) wherein R^{20} is of the formula $-N(CO_2R^{27})CO(1-$
10 4C)alkyl is conveniently prepared by reacting a compound of the formula (I) and (II) wherein R^1 or R^{20} is hydroxy with an amide of the formula $HN(CO_2R^{27})CO(1-4C)alkyl$ under Mitsunobu conditions. For example, in the presence of tri-n-butylphosphine and 1,1'-(azodicarbonyl)dipiperidine in an organic solvent such as THF, and in the temperature range 0°C - 60°C, but preferably at ambient temperature. Details of analogous Mitsunobu reactions
15 are contained in Tsunoda et al, Tet. Letts., 34, 1639, (1993). Amides of the formula $HN(CO_2R^{27})CO(1-4C)alkyl$ may be prepared by standard procedures of organic chemistry which are within the ordinary skill of an organic chemist.

A compound of the formula (I) or (II) wherein R^1 or R^{20} is chloro may also be prepared from a compound of the formula (XIII), by reacting the latter compound with lithium
20 chloride and crown ether, in a suitable organic solvent such as THF, in a temperature range of ambient temperature to reflux. A compound of the formula (I) or (II) wherein R^1 or R^{20} is (1-4C)alkylthio or (1-4C)alkoxy may be prepared by reacting the compound of the formula (XIII) with sodium thio(1-4C)alkoxide or sodium (1-4C)alkoxide respectively, in an alcohol or THF, in a temperature range of 0°C to reflux.

25 It is also possible to convert one R^7 group into another R^7 group as a final step in the preparation of a compound of the formula (I) or (II) (see the specific examples).

When an optically active form of a compound of the formula (I) is required, it may be obtained by carrying out one of the above procedures using an optically active starting material, or by resolution of a racemic form of the compound or intermediate using a standard
30 procedure.

Similarly, when a pure regioisomer of a compound of the formula (I) is required, it may be obtained by carrying out one of the above procedures using a pure regioisomer as a starting material, or by resolution of a mixture of the regioisomers or intermediates using a standard procedure.

5 According to a further feature of the invention there is provided a compound of the formula (I), or a pharmaceutically-acceptable salt thereof, for use in a method of treatment of the human or animal body by therapy.

According to a further feature of the present invention there is provided a method for producing an antibacterial effect in a warm blooded animal, such as man, in need of such
10 treatment, which comprises administering to said animal an effective amount of a compound of the present invention, or a pharmaceutically-acceptable salt thereof.

The invention also provides a compound of the formula (I), or a pharmaceutically-acceptable salt thereof, for use as a medicament; and the use of a compound of the formula (I) of the present invention, or a pharmaceutically-acceptable salt thereof, in the manufacture of a
15 novel medicament for use in the production of an antibacterial effect in a warm blooded animal, such as man.

In order to use a compound of the formula (I) or a pharmaceutically-acceptable salt thereof for the therapeutic treatment of mammals including humans, in particular in treating infection, it is normally formulated in accordance with standard pharmaceutical practice as a
20 pharmaceutical composition.

Therefore in another aspect the present invention provides a pharmaceutical composition which comprises a compound of the formula (I) or a pharmaceutically-acceptable salt thereof and a pharmaceutically-acceptable diluent or carrier.

The pharmaceutical compositions of this invention may be administered in standard
25 manner for the disease condition that it is desired to treat, for example by oral, rectal or parenteral administration. For these purposes the compounds of this invention may be formulated by means known in the art into the form of, for example, tablets, capsules, aqueous or oily solutions or suspensions, (lipid) emulsions, dispersible powders, suppositories, ointments, creams, drops and sterile injectable aqueous or oily solutions or suspensions.

30 In addition to the compounds of the present invention the pharmaceutical composition of this invention may also contain or be co-administered with one or more known

drugs selected from other clinically useful antibacterial agents (for example β -lactams or aminoglycosides). These may include penicillins, for example oxacillin or flucloxacillin and carbapenems, for example meropenem or imipenem, to broaden the therapeutic effectiveness against methicillin-resistant staphylococci. Compounds of this invention may also contain or
5 be co-administered with bactericidal/permeability-increasing protein product (BPI) or efflux pump inhibitors to improve activity against gram negative bacteria and bacteria resistant to antimicrobial agents.

A suitable pharmaceutical composition of this invention is one suitable for oral administration in unit dosage form, for example a tablet or capsule which contains between
10 100mg and 1g of the compound of this invention.

In another aspect a pharmaceutical composition of the invention is one suitable for intravenous, subcutaneous or intramuscular injection.

Each patient may receive, for example, a daily intravenous, subcutaneous or intramuscular dose of 5 mgkg^{-1} to 20 mgkg^{-1} of the compound of this invention, the
15 composition being administered 1 to 4 times per day. The intravenous, subcutaneous and intramuscular dose may be given by means of a bolus injection. Alternatively the intravenous dose may be given by continuous infusion over a period of time. Alternatively each patient will receive a daily oral dose which is approximately equivalent to the daily parenteral dose, the composition being administered 1 to 4 times per day.

20

Antibacterial Activity

The pharmaceutically-acceptable compounds of the present invention are useful antibacterial agents having a good spectrum of activity in vitro against standard Gram-positive organisms, which are used to screen for activity against pathogenic bacteria. Notably, the
25 pharmaceutically-acceptable compounds of the present invention show activity against enterococci, pneumococci and methicillin resistant strains of *S. aureus* and coagulase negative staphylococci. The antibacterial spectrum and potency of a particular compound may be determined in a standard test system.

The antibacterial properties of the compounds of the invention may also be
30 demonstrated in-vivo in conventional tests. No overt toxicity or other untoward effects are observed when compounds of the formula I are so tested.

The following results were obtained on a standard in-vitro test system. The activity is described in terms of the minimum inhibitory concentration (MIC) determined by the agar-dilution technique with an inoculum size of 10^4 CFU/spot.

Staphylococci were tested on agar, using an inoculum of 10^4 CFU/spot and an incubation temperature of 37°C for 24 hours - standard test conditions for the expression of methicillin resistance.

Streptococci and enterococci were tested on agar supplemented with 5% defibrinated horse blood, an inoculum of 10^4 CFU/spot and an incubation temperature of 37°C in an atmosphere of 5% carbon dioxide for 48 hours - blood is required for the growth of some of the test organisms.

<u>Organism</u>	<u>MIC ($\mu\text{g/ml}$)</u>
<u>Example 1</u>	
Staphylococcus aureus:	
Oxford	1.0
15	Novb. Res
	0.5
	MRQS
	0.5
	MRQR
	0.5
Coagulase Negative Staphylococcus	
	MS
	0.25
20	MR
	1.0
Streptococcus pyogenes	
	C203
	1.0
Enterococcus faecalis	
	1.0
Bacillus subtilis	
	0.25

25

Novb. Res = Novobiocin resistant

MRQS = methicillin resistant quinolone sensitive

MRQR = methicillin resistant quinolone resistant

MR = methicillin resistant

30 MS = methicillin sensitive

The invention is now illustrated but not limited by the following Examples in which unless otherwise stated :-

- i) evaporations were carried out by rotary evaporation in vacuo and work-up
5 procedures were carried out after removal of residual solids by filtration;
- (ii) operations were carried out at ambient temperature, that is in the range 18-26°C and in air unless otherwise stated, or unless the skilled person would otherwise work under an inert atmosphere;
- (iii) column chromatography (by the flash procedure) was performed on Merck
10 Kieselgel silica (Art. 9385) unless otherwise stated;
- (iv) yields are given for illustration only and are not necessarily the maximum attainable;
- (v) the end-products of the formula I generally have satisfactory microanalyses and their structures were confirmed by NMR and mass spectral techniques [proton magnetic
15 resonance spectra were determined in DMSO-D₆ unless otherwise stated using a Varian Gemini 2000 spectrometer operating at a field strength of 300 MHz, or a Bruker AM250 spectrometer operating at a field strength of 250 MHz; chemical shifts are reported in parts per million downfield from tetramethylsilane as an internal standard (δ scale) and peak multiplicities are shown thus: s, singlet; d, doublet; AB or dd, doublet of doublets; t, triplet.
20 m, multiplet; fast-atom bombardment (FAB) mass spectral data were obtained using a Platform spectrometer (supplied by Micromass) run in electrospray and, where appropriate, either positive ion data or negative ion data were collected];
- (vi) intermediates were not generally fully characterised and purity was in general assessed by thin layer chromatographic, infra-red (IR), mass spectral (MS) or NMR analysis;
25 and
- (vii) in which :-

	MPLC	is medium pressure chromatography
	TLC	is thin layer chromatography
30	DMSO	is dimethylsulfoxide
	CDCl ₃	is deuterated chloroform

	MS	is mass spectroscopy
	ESP	is electrospray
	CI	is chemical ionization
	DMF	is N,N-dimethylformamide
5	THF	is tetrahydrofuran
	LDA	is lithium diisopropylamide
	TFA	is trifluoroacetic acid
	NMP	is N-methylpyrrolidone
	dba	is dibenzylideneacetone

10

Reference Example 1 : (5S)-Acetamidomethyl-3-(4-trimethyltinphenyl)oxazolidin-2-one

Hexamethyldistannane (1.77g) followed by Pd(PPh₃)Cl₂ (155mg) was added to a partial solution of 5S-acetamidomethyl-3-(4-iodophenyl)oxazolidin-2-one (1.84g) in dioxan (25ml). The vessel was purged well with argon and the reaction mixture was stirred at 95-100°C for 6
15 hours, after which the reaction was judged as complete by TLC. The solution was decanted from a film of black gum which had separated, and evaporated. The title product was isolated as an oil by MPLC (Merck 9385 silica using as eluant a mixture of methanol and dichloromethane increasing in polarity from 5% to 10% methanol) which crystallised in 59% yield (1.19 g) on trituration with ether.

20

See also WO Patent 94-13649 (23.06.94) for preparation of Reference Example 1.

NMR (300MHz, DMSO-D₆): δ 0.25(s,9H), 1.84(s,3H), 3.42(t,2H), 3.74(d of d,1H), 4.10(t,1H), 7.50(m,4H), 4.70(m,1H), 8.22(t,1H).

25 MS: ESP+ (M+H)=397.

Reference Example 2 : N-((5S)-3-(3-Fluoro-4-(1-tert-butoxycarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Lithium chloride (1.53g) was added to a stirred solution of of t-butyl-1,2,3,6-tetrahydro-4-
30 (trifluoromethylsulfonyloxy)pyridine-1-carboxylate (4.97g) in NMP (50ml), followed by Pd₂(dba)₃ (550mg) and the vessel was purged well with argon. After stirring for 5 minutes at

ambient temperature, a solution of N-((5S)-3-(3-fluoro-4-trimethyltinphenyl)-2-oxooxazolidin-5-ylmethyl)acetamide (4.99g, WO Patent 94-13649, 23.06.94) in NMP (20ml) was added. The reaction mixture was stirred at ambient temperature for 5 hours, then at 50°C for 18 hours. TLC (ethyl acetate) indicated that the reaction was complete.

- 5 A 2.0M aqueous potassium fluoride solution (12ml) was added and the reaction mixture stirred at ambient temperature for 1½ hours. Water was added and the product was extracted with ethyl acetate. The organic phase was washed with water, brine, dried over anhydrous sodium sulfate and evaporated to a gum. The title compound was isolated as a gel in 68% yield (4.39g) by MPLC (Merck 9385 silica, using as eluant a mixture of methanol (3.5%)
10 and dichloromethane), and used in subsequent steps without further purification.

NMR (300MHz, DMSO-D6): δ 2.40(m, 1H), 3.40(m, 3H), 3.50(m, 1H), 3.70(m, 1H), 3.98(m, 2H), 4.10(m, 1H), 4.70(m, 1H), 5.95(m, 1H), 7.35(m, 3H), 8.20(t, 1H).

Reference Example 3 : 5,6-Dihydro-4-trifluoromethylsulfonyloxy-2H-pyran

- 15 LDA / THF (31.5ml of 1.92M solution) was slowly added to a stirred solution of tetrahydro-4H-pyran-4-one (5.5g) in THF (30ml) at -70°C, under argon. The mixture was stirred for 30 minutes at -70°C and then a solution of N-phenyl triflimide (21.6g) in THF (30mls) was added. The mixture was allowed to warm to ambient temperature and stirred for 18 hours. The reaction mixture was evaporated and subjected to chromatography by MPLC on Alumina
20 (ICN, N32-63, using as eluant a mixture of ethyl acetate (5%) and iso-hexane). The product was distilled by Kugelrohr (100°C / 10mm). Remaining traces of the triflimide reagent were removed by a second MPLC (Silica, using as eluant a mixture of ethyl acetate (5%) and iso-hexane) followed by a second Kugelrohr distillation, giving the title compound as a colourless oil in 40% yield (5.1g), which was stored at -20°C.
25 NMR (300MHz, CDCl₃): δ 2.24(m,2H), 3.90(m,2H), 4.25(m,2H), 5.82(m,1H).

Reference Example 4 : 1-(4-Bromo-3-fluorophenyl)-2,2,5,5-tetramethyl-1,2,5-azadisilalidine

- Triethylamine (56.8g) was added to a stirred solution of 4-bromo-3-fluoroaniline(42.75g, 30 JCS, 2815 (1958)) in dichloromethane (400ml) followed by the slow addition, with ice-bath cooling, of a solution of 1,2-bis(chlorodimethylsilyl)ethane (50.79g) in dichloromethane

(100ml). After stirring for 2 days, an NMR of a worked up sample showed that the reaction was complete. The reaction mixture was washed with ice cold 2N aqueous Na_2HPO_4 and water, dried over anhydrous magnesium sulfate and evaporated to give a brown oil. The title compound was isolated in 93% yield (69.4g) by vacuum distillation. (bp = 105-110°C / 0.2bar).

NMR (300MHz CDCl_3): δ 0.22 (s, 12H), 0.85 (s, 8H), 6.52 (d of d, 1H), 6.65 (d of d, 1H), 7.30 (t, 1H).

Reference Example 5 : 4-(1-Benzyl-4-hydroxy-4-piperidyl)-3-fluoroaniline

10 n-Butyllithium/hexane (129ml of 1.40M) was slowly added to a stirred solution of Reference Example 4 (59.76g) in dry THF (300ml) at -70°C under argon. The solution was stirred for 5 minutes, then a solution of N-benzyl-4-piperidone (37.8g) in THF (150ml) was slowly added. The reaction mixture was allowed to warm to ambient temperature and stirred for 3 hours. The reaction mixture was then quenched by the addition of ice, followed by water, then
15 acidified to <pH1 by the addition of 5N aqueous HCl. After stirring for 5 minutes the mixture was extracted with ether and the organic phase discarded. The aqueous phase was basified with a slight excess of aqueous sodium hydroxide and the product was extracted with ether. The organic phase was washed with brine, dried over anhydrous sodium sulfate and evaporated to give the title compound as an oil in which crystallized on trituration with
20 cyclohexane (yield = 39.49g, 73%)

NMR (300MHz $\text{DMSO}-d_6$): δ 1.55(d, 2H), 2.10(d of t, 2H), 2.42(m, 2H), 2.55(obsured by DMSO), 3.50(s, 2H), 4.65(s, 1H), 5.18(s, 2H), 6.25(d of d, 1H), 6.33(d of d, 1H), 7.30(m, 6H).

MS: ESP+ (M+H) = 301.

25

Reference Example 6 : 3-Fluoro-4-(1-benzyl-1,2,5,6-tetrahydropyrid-4-yl)aniline

A solution of Reference Example 5 (36.0g) in 5N aqueous HCl (200ml) was heated under reflux for 9 hours. Concentrated HCl (50ml) was added and reflux was continued for a further 4 hours. The reaction mixture was basified, with cooling, with 40% aqueous sodium
30 hydroxide solution and extracted with ether. Addition of aqueous ammonium hydroxide solution avoided problems with insoluble partial salts. The organic phase was washed with

brine, dried over anhydrous sodium sulfate and evaporated well giving the title compound as a viscous gum in 96% yield (32.33g).

NMR (300MHz DMSO-D6): δ 2.35(s, 2H), 2.55(t, 2H), 2.98(d, 2H), 3.53(s, 2H), 5.35(s, 2H), 5.75(s, 1H), 6.29(m, 2H), 6.95(t, 1H), 7.30(m, 5H).

5 MS: ESP+ (M+H) = 283.

Reference Example 7 : N-Benzylloxycarbonyl-3-fluoro-4-(1-benzyl-1,2,5,6-tetrahydropyrid-4-yl)aniline

Dibenzyl dicarbonate (43g) was slowly added to a stirred solution of Reference Example 6
10 (28.2g) in dichloromethane (200ml) at 0-5°C. The reaction mixture was allowed to warm to ambient temperature and stirred for 18 hours. The solution was evaporated to an oil which was triturated with ether. Sticky, insoluble material was discarded and the filtrate was purified by chromatography (MPLC on silica, using as eluant a mixture of ethyl acetate (25%) and iso-hexane). The product was taken into ethyl acetate and precipitated as the HCl salt by addition
15 of a mixture of HCl and ethyl acetate. The resulting product was filtered off, washed with ethyl acetate and dried under vacuum to give the title product.

Yield = 10.2g (23%)

NMR (300MHz DMSO-D6): δ 2.61(m, 1H), 2.85(m, 1H), 3.20(m, 1H), 3.53(m, 1H), 3.75(s, 2H), 4.40(t, 2H), 5.15(s, 2H), 5.95(s, 1H), 7.35(m, 11H), 7.60(m, 2H).

20 MS: ESP+ (M+H) = 417.

Reference Example 8 : N-tert-Butoxycarbonyl-N-((5R)-3-(3-fluoro-4-(N-benzyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

BOC-acetamide (2.85g, Chem.Pharm.Bull.Jap.,36, 3125 (1988)) was added to a stirred partial
25 solution of Example 13 (4.55g) in dry THF (80ml) and the mixture cooled to 0-4°C under argon. Tributylphosphine (3.62g) was added, followed by portion-wise addition of 1,1'-(azodicarbonyl)-dipiperidine (4.51g). The reaction mixture was stirred for 30 minutes at 0-4°C during which time a precipitate formed. The reaction mixture was allowed to warm to ambient temperature and stirred for 18 hours. The reaction mixture was filtered and
30 evaporated to a gum. The title compound was rapidly isolated by MPLC (silica, using as

eluant a mixture of ethyl acetate (70%) and iso-hexane) as a gum and subjected to the minimum of handling (brief evaporation at <40°C) before deprotection in Example 12.

Yield = 6.9g (inc. some solvent).

5 **Reference Example 9: 4-(1-tert-butyloxycarbonyl-1,2,5,6-tetrahydropyrid-4-yl)-3,5-difluoroaniline.**

nBuLi (1.6M in hexanes, 33.3ml) was added to a solution of N,N-[1,2-bis(dimethylsilyl)ethane]-3,5-difluoroaniline (12.0g, J.Org. Chem., 60, 5255-5261 (1995)) in 150ml THF at -70°C, maintaining temperature below -70°C during the addition. The mixture
10 was then stirred at that temperature for 4 hours. A solution of ZnCl₂ (0.5M in THF, 106.7ml) was then added at < 60°C, the mixture allowed to warm to ambient temperature and then degassed by bubbling argon through the solution for 15 minutes. Tetrakis(triphenylphosphine) Pd⁰ (1.54g) was then added together with t-butyl-1,2,5,6-tetrahydro-4-(trifluoromethylsulfonyloxy)-pyridine carboxylate (15.4g, Synthesis, 993 (1991)), the mixture
15 degassed again and stirred overnight at ambient temperature. After refluxing for two hours the mixture was cooled to 10°C and added to 100ml ice water. After adjusting the pH to 2-3 with 5N HCl and stirring at 10°C for 5 minutes the mixture was brought to pH 8 with NaHCO₃, extracted into ethyl acetate (2x150ml) and evaporated to give a dark oil which was chromatographed on alumina (ICN-N-32-63) with isohexane/ethyl acetate 4:1. Trituration of
20 the resulting oil gave a solid (6.62g, 48%).

NMR (300MHz, DMSO-D₆) d: 1.46(s, 9H), 2.27(broad s, 2H), 3.52(d/d, 2H), 3.97(broad s, 2H), 5.70(s, 3H), 6.20(d, 2H).

25 **Reference Example 10: N-Benzylloxycarbonyl-4-(1-tert-butyloxycarbonyl-1,2,5,6-tetrahydropyrid-4-yl)-3,5-difluoroaniline.**

NaHCO₃ (5.5g) was added to a mixture of Reference Example 9 (10.0g), acetone (150ml) and water (75ml) and the mixture cooled in an ice-bath. A solution of benzylchloroformate (5.3ml) in 10ml of acetone was added slowly, the mixture was stirred whilst the temperature was warmed to ambient and then stirred for a further hour. Water (100ml) was added and the
30 mixture extracted with ethyl acetate (100ml + 2x50ml). The combined extracts were dried

and evaporated to give a solid which was recrystallised from 90ml of acetonitrile to give a solid (11.2g). Recrystallisation of the residues gave further solid (0.73g). Total yield (83%).

NMR (300MHz, DMSO-D6) d : 1.42(s, 9H), 2.25(broad s, 2H), 3.50(t, 2H), 3.95(broad s, 2H), 5.15(s, 2H), 5.78(s, 1H), 7.15(d, 2H), 7.38(m, 5H), 10.16(broad s, 1H).

5

Reference Example 11: (5R)-Hydroxymethyl-3-(3,5-difluoro-4-(tert-butyloxycarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)oxazolidin-2-one.

nBuLi (1.45M in hexanes, 15.7ml) was added dropwise at < -70°C to a solution of Reference Example 10 (9.65g) in anhydrous THF (200ml) at -70°C. After 20 minutes a solution of (R)-glycidyl butyrate (3.44g) in 25ml THF was added at the same temperature. After the addition the cooling bath was removed and the reaction allowed to stir to ambient temperature overnight. After addition of methanol (15ml) and stirring for 10 minutes, 100ml ethyl acetate was added, the mixture cooled to 10°C and 0.5M HCl (20ml) added slowly. After separation of the aqueous phase it was re-extracted with ethyl acetate (2x100ml) and the combined organic phases were washed with saturated NaHCO₃ solution, with brine and evaporated to give an oil (12.0g) . Chromatography on silica gel with ethyl acetate gave a solid (7.66g, 86%).

NMR (300MHz, DMSO-D6) d : 1.40(s, 9H), 2.27(broad s, 2H), 3.51(m, 3H), 3.64/3.68(m/m, 1H), 3.80(m, 1H), 3.97(broad s, 2H), 4.05(t, 1H), 4.70(m, 1H), 5.18(t, 1H), 5.82(broad s, 1H), 7.32(m, 2H).

Reference Example 12: 3,5-difluoro-4-(2,3-dihydropyran-4-yl)aniline

nBuLi (1.4M in hexanes, 63.5ml) was added dropwise to a solution of N,N-[1,2-bis(dimethylsilyl)ethane]-3,5-difluoranyl (20.0g, J.Org.Chem., 60, 5255-5261 (1995)) in 200ml THF at -70°C. The mixture was stirred at that temperature for 4 hours. Chlorotitanium triisopropoxide (1.0M in hexanes, 89ml) was then added dropwise at the same temperature, and the mixture allowed to warm to ambient temperature for 30minutes. The mixture was recooled to -70°C and tetrahydro-4H-pyran-4-one (9.0g in a few ml of THF) added dropwise and the reaction mixture allowed to stir at ambient temperature for 48 hours. After cooling in an ice bath, conc.HCl was added to pH 1-2, the organic phase separated and the aqueous phase reextracted with ether. The aqueous phase was adjusted to pH 9 with 2.5N NaOH, a

small amount of solid filtered off and the filtrate extracted with ethyl acetate (4x100ml).

Work-up (ie. washing and evaporation of solvent) gave an oil which was dissolved in 75ml dioxan and 75ml 5N HCl. The mixture was refluxed for 15 minutes and the dioxan removed in vacuo. The residue was retaken up in dilute HCl, extracted with ether. The aqueous layer
5 was basified to pH 8 with NaHCO₃ and extracted with ethyl acetate (3x75ml). Evaporation in vacuo gave a solid (8.5g, 45%).

NMR (300MHz, DMSO-D₆) d : 2.20(broad s, 2H), 3.74(t, 2H), 4.12(q, 2H), 5.65(s, 2H), 5.70(s, 1H), 6.10(s, 1H), 6.18(s, H).

MS: ESP+ (M+H) = 212.

10

Reference Example 13: N-Benzylloxycarbonyl-4-(2,3-dihydro-6H-pyran-4-yl)-3,5-difluoroaniline

Benzylchloroformate (6.6ml, 7.88g) was added to a mixture of Reference Example 12 (8.4g), acetone (150ml) and water (75ml) containing NaHCO₃ (7g) in an ice-bath. The mixture was
15 stirred for 3 hours whilst warming to ambient temperature. At the end of this period 100ml water was added, the mixture extracted with ethyl acetate (3x100ml) and evaporated in vacuo to give a crude solid which was triturated with ether to give a solid (75%).

NMR (300MHz, DMSO-D₆) d : 2.25(broad s, 2H), 3.76(t, 2H), 4.18(m, 2H), 5.15(s, 2H), 5.85(s, 1H), 7.14(s, 1H), 7.18(s, 1H), 7.38(m, 5H), 10.16(broad s, 1H).

20

Reference Example 14 : (5R)-Hydroxymethyl-3-(3,5-difluoro-4-(2,3-dihydro-6H-pyran-4-yl)phenyl)oxazolidin-2-one

nBuLi (1.4M in hexanes, 23.2ml) was added dropwise to a solution of Reference Example 13 (10.2g) in THF (200ml) at -70°C and the mixture stirred for 20 minutes. A solution of (R)-
25 glycidyl butyrate (4.68g) in THF (20ml) was added dropwise and the reaction allowed to stir over 48 hours, warming to ambient temperature over a few hours. Methanol (20ml) was added, the mixture stirred for 10 minutes, then ethyl acetate (100ml) followed by 100ml 0.5N HCl added. The organic phase was separated, the aqueous re-extracted and the combined organic phases washed with saturated NaHCO₃ solution, followed by evaporation in vacuo to
30 give a crude solid which was purified on silica-gel using ethyl acetate as eluant to give a solid (7.4g, 80%).

NMR (300MHz, DMSO-D6) d: 2.29(broad s, 2H), 3.53(m, 1H), 3.67(m, 1H), 3.79(m, 3H), 4.06(t, 1H), 4.19(m, 2H), 4.71(m, 1H), 5.19(t, 1H), 5.89(m, 1H), 7.30(s, 1H), 7.36(s, 1H).

MS: ESP+ (M+H) = 312.

5 **Reference Example 15: (2RS)-2-Methyl-4-trifluoromethylsulfonyloxy-3,6- and -5,6-dihydropyran.**

(RS)-2-Methyl-tetrahydro-4H-pyran-4-one (1.04 g, Chem.Ber., 88, 1053, (1955)) in dry THF (30 ml) at -78 °C was treated with LDA (2.0 M in heptane/THF/PhEt) (4.56 ml) under argon. After 45 minutes *N*-phenyl-bis(trifluoromethanesulfonimide) (3.58 g) was added in one go
10 and stirring was continued for 1 hour at -78°C. The ice bath was then removed and stirring continued for a further 19 hours. The solution was evaporated and the residue purified by firstly silica-gel MPLC [using a mixture of 4% ethyl acetate and hexanes as eluant], and then by bulb-to-bulb distillation. Vinyl trifluoromethanesulfonate (1.36g, 60%) as a 1.4 : 1 mixture of the title methyl-regioisomers was collected as the fraction at 120°C/10 mmHg.
15 NMR (250MHz, CDCl₃) d: 1.29 and 1.31 (m/m, 3H each), 2.17-2.43 and 2.54-2.70 (m/m, 2H each), 3.66-3.82 (m, 2H), 4.10 (ddd, 1H), 4.19-4.40 (m, 3H), 5.68-5.74 (m, 1H) and 5.79-5.84 (m, 1H).

Reference Example 16: 2,6-Dimethyl-4-hydroxy-2,3,5,6-tetrahydro-4H-pyran.

20 2,6-Dimethyl-4H-pyran-4-one (5.0 g) in ethanol (40 ml) was hydrogenated under 10% Pd-C (0.5 g) at 10 bar of hydrogen for 2 days. The solution was then evaporated to an oil which was purified by silica-gel MPLC [using a 1:1 mixture of ethyl acetate and hexane as eluant] to give the title product (3.71g, 74%).

NMR (300 MHz, CDCl₃) d: 1.16-1.25 (m, 6H), 1.40-1.55 (m, 2H), 1.60-1.69 (m, 1H),
25 1.89-1.99 (m, 1H), 3.40-3.50 (m, 1H), 3.70-3.99 and 4.21 (m/m, together 2H).

Reference Example 17: 2,6-Dimethyl-2,3,5,6-tetrahydro-4H-pyran-4-one

Reference Example 16 (3.71 g) was dissolved in dry dichloromethane (50ml) and pyridine chlorochromate (PCC) (9.2 g) was added portionwise with stirring. The mixture was stirred
30 for 18 hours, filtered through Celite and evaporated. The residue was purified by bulb-to-bulb distillation (90°C, 16 mbar) to afford the title ketone (1.09g, 30%).

NMR (300 MHz, CDCl₃) d: 1.32 (d, 6H), 2.17-2.40 (m, 4H) and 3.68-3.80 (m, 2H).

Reference Example 18: 2,6-Dimethyl-4-trifluoromethylsulfonyloxy-5,6-dihdropyran.

Reference Example 17 (1.09 g) in dry THF (10 ml) at -78°C was treated with LDA (2.0 M in 5 heptane/THF/PhEt) (5.55 ml) under argon. After 30 minutes *N*-phenyl-bis(trifluoromethanesulfonylimide) (3.21 g) was added in one go and stirring was continued for 1 hour at -78°C. The ice bath was then removed and stirring continued for a further 19 hours. The solution was evaporated and the residue purified by silica-gel MPLC [using a mixture of 4% ethyl acetate and hexanes as eluant] to give the vinyl triflate (1.75g, 79%).

10 NMR (300 MHz, CDCl₃) d: 1.50 and 1.51 (d/d, each 3H), 2.19-2.40 (m, 2H), 3.70-3.83 (m, 1H), 4.30-4.40 (m, 1H) and 5.70 (s, 1H).

Reference Example 19: There is no compound with Reference Example No. 19.

Reference Example 20: 2-Benzylloxymethyl-2,3,5,6-tetrahydro-4H-pyran-4-one.

A mixture of (RS)-2-benzylloxymethyl-2,3-dihydro-4H-pyran-4-one (0.82g, J. Org. Chem., 15 44, 811 (1979)) in ethanol (15 ml) containing 10% Pd-C (0.089 g) and triethylamine (10 drops) was hydrogenated until NMR spectroscopy indicated almost complete reaction. The solution was evaporated and the residue purified by silica-gel MPLC [25% EtOAc/hexanes] and then bulb-to-bulb distillation [220 °C/2 mmHg] to give the title ketone (0.47g, 56%).

NMR (300 MHz, CDCl₃) d: 2.30-2.40 (m, 2H), 2.21-2.70 (m, 2H), 3.55 (d, 2H), 3.70 (dt, 20 1H), 3.80-3.89 (m, 1H), 4.35 (dd, 1H), 4.60 (s, 2H) and 7.30-7.39 (m, 5H).

MS: ESP+ (M+H) = 221.

Reference Example 21: (2RS)-2-Benzylloxymethyl-4-trifluoromethylsulfonyloxy-3,6- and -5,6-dihdropyran.

Under analogous conditions to those described in Reference Example 18, Reference Example 25 20 (6.1 g) was transformed into the crude vinyltriflate. Silica-gel MPLC [using a mixture of 10% ethyl acetate and hexanes as eluant], then alumina MPLC [using a mixture of 5% ethyl acetate and hexanes as eluant] gave a mixture of the title products (5.9 g, 61%).

NMR (300 MHz, CDCl₃) d : 2.20-2.31 (m, 2H), 2.42-2.70 (m, 2H), 3.45-3.62 (m, 4H), 3.70-3.90 (m, 2H), 4.10-4.19 (m, 1H), 4.22-4.48 (m, 3H), 4.52-4.63 (m, 4H), 5.79 (s, 1H), 5.80-5.82 (m, 1H) and 7.25-7.40 (m, 10H).

5 Reference Example 22: N-((5S)-3-(4-(2-methanesulfonyloxymethyl-3,6- and -5,6-dihydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

Mesyl chloride (0.080 g, 0.054 ml) was added slowly to a stirred mixture of Example 52 (0.183 g), triethylamine (0.066 g, 0.090 ml) and dichloromethane (8 ml) at 0°C. After 1 hour, more mesyl chloride (0.070 ml) and triethylamine (0.070 ml) were added and stirring
10 continued for a further 2 hours. The solution was then washed with 10% HCl, saturated aqueous NaHCO₃ solution and with water, then dried and evaporated to an oil which was purified by Isolute silica-gel chromatography [using a mixture of 6% MeOH and CH₂Cl₂ as eluant] to give a mixture of the title mesylates (0.190g, 85%).

NMR (300 MHz, CDCl₃) d : 2.05 (s, 3H), 2.31-2.55 (m, 1H), 2.91-3.05 (m, 1H m), 3.06, 3.08
15 and 3.12 (s/s/s, together 3H), 3.60-4.28 (m, 6H), 4.30-4.64 (m, 3H), 4.75-4.90 (m, 1H), 5.96, 6.12 and 6.30 (each broad s, together 2H) and 7.32-7.60 (m, 4 H).

MS: ESP+ (M+H) = 425.

Reference Example 23: N-((5S)-3-(4-(2-azidomethyl-3,6- and 5,6-dihydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

20 The mixture of mesylates Reference Example 22 (0.177 g) and sodium azide (0.041 g) in dimethyl sulfoxide (2 ml) were stirred at 80°C for 5 hours. NMR indicated very little consumption of starting material and more sodium azide (0.050 g) was added. The mixture was heated at 80°C for 18 hours. The resulting mixture was dissolved in ethyl acetate and washed twice with water. The organic layer was dried and evaporated to afford the title
25 mixture of azides (0.138g, 89%).

NMR (300 MHz, CDCl₃) d : 2.01 and 2.05 (s/s, together 3H), 2.28-2.53 (m, 2H), 3.38-3.49 (mm, 2H), 3.60-3.70 (m, 2H), 3.78-3.90 (m, 2H), 4.09 (t, 2H), 4.40-4.50 (m, 1H), 4.72-4.95 (m, 1H), 5.98 and 6.12 (s/s, together 1H), 6.50 (broad t, 1H,) and 7.30-7.55 (m, 4H).

MS: ESP+ (M+H) = 372.

Reference Example 24: (2RS)-2-Benzylloxymethyl-4-trifluoromethylsulfonyloxy-3,6-dihydropyran.

(RS)-2-Benzylloxymethyl-2,3-dihydro-4H-pyran-4-one (5.69g, J. Org. Chem., 44, 811
5 (1979)) in dry THF (200 ml) was treated with L-selectride (27ml, 1.0M in THF) at -78°C and
stirred for 1.25 hour. N-Phenyl-bis(trifluoromethanesulfonimide) (10.35 g) was added and
stirring continued for 3 days with warming to ambient temperature. The solution was
evaporated to an oil which was purified by firstly silica-gel MPLC [10% EtOAc/hexanes as
eluant], then alumina MPLC [10% EtOAc/hexanes as eluant] to afford the title vinyl triflates
10 (6.00g, 66%).

NMR (300MHz CDCl₃) d: 2.23 (dt, 1H), 2.41-2.58 (m, 1H), 3.50-3.62 (m, 2H), 3.80-3.90 (m,
1H), 4.22-4.43 (m, 2H), 4.60 (s, 2H), 5.80 (d, 1H), 7.25-7.41 (m, 5H).

Example 1 : N-((5S)-3-(4-(1-tert-butoxycarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxo-oxazolidin-5-ylmethyl)acetamide

To a stirred solution of t-butyl-1,2,3,6-tetrahydro-4-(trifluoromethylsulfonyloxy)-pyridine-1-
carboxylate (497mg, Synthesis, 993 (1991)) in NMP (8ml), was added Pd₂(dba)₃ (60mg),
triphenylarsine (80mg) and lithium chloride (166mg), and the vessel was purged well with
argon. After stirring for 5 mins. at ambient temperature, a solution of the starting material
20 (516mg, Reference Example 1) in NMP (2ml) was added and the reaction mixture stirred at
ambient temperature for 18hours. TLC (ethyl acetate) indicated incomplete reaction so more t-
butyl-1,2,3,6-tetrahydro-4-(trifluoromethylsulfonyloxy)pyridine-1-carboxylate (497mg) was
added and the reaction mixture was heated at 40°C for a further 24hours, after which the
reaction was complete. A 1.0M solution of aqueous potassium fluoride (1.0ml) was added and
25 the reaction mixture was stirred at ambient temperature for 30 mins. Water was added and the
product was extracted with ethyl acetate, with filtration of the 2 phase mixture. The organic
phase was washed with water, saturated aqueous sodium chloride solution, dried over
anhydrous sodium sulfate and evaporated to a semi-solid gum. The product was isolated by
MPLC (Merck 9385 silica, using as eluant a mixture of acetonitrile and ethyl acetate

increasing in polarity from 0% to 50% acetonitrile over 15 minutes), and was crystallised by dissolving in a small volume of ethyl acetate and triturating with ether (yield = 125mg, 23%).

NMR (200MHz, DMSO-D6): δ 1.42(s,9H), 2.43(m, obscured by DMSO), 3.42(t,2H), 3.52(t,2H), 3.75(d of d,1H), 3.98(d,2H), 4.16(t,1H), 4.71(m,1H), 6.12(s,1H), 7.49(AB,4H),
5 8.20(t,1H).

MS: ESP+ (M+H) = 416.

Example 2 : N-((5S)-3-(4-(1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

10 A general procedure for deprotection was used as follows. The starting material (Example 1) was dissolved in a small volume of trifluoroacetic acid and the yellow solution was heated briefly to reflux. Ether (10x volume of TFA used) was added, precipitating a sticky gum. The solvent was decanted off and the gum crystallised on trituration with ether. The TFA salt of the product was filtered off, washed with ether and dried under vacuum.

15 For example, 10.5 mg of starting material in 0.1ml of TFA was treated in this way to give the title compound in 97% yield (10.5mg).

NMR (300MHz, DMSO-D6): δ 1.81(s,3H), 2.65(s,2H), 3.35(partially obscured by H₂O), 3.42(t,2H), 3.75(m,3H), 4.13(t,1H), 4.73(m,1H), 6.18(s,1H), 7.54(AB,4H), 8.25(t,1H),
8.85(s,2H).

20 MS: ESP+ (M+H) = 316.

Example 3 : N-((5S)-3-(4-(1-methoxycarbonyl-1,2,5,6-tetrahydropyrid-4-yl))phenyl)-2-oxo-oxazolidin-5-ylmethyl)acetamide

To a stirred solution of the starting material (Example 2) (129mg) in a mixture of acetone
25 (2ml) and water (1ml) was added sodium hydrogen carbonate (101mg), and the mixture was cooled to 0-5°C. Methyl chloroformate (34mg, 28 μ l) was added and stirring continued at 0-5°C for 30 minutes. Then the reaction mixture was allowed to warm to ambient temperature over 30 minutes, after which TLC (10% methanol in dichloromethane) showed the reaction was complete. The reaction mixture was diluted with water and extracted with ethyl acetate.
30 The organic phase was washed with saturated aqueous sodium chloride solution. dried over

anhydrous sodium sulfate and evaporated to give a crystalline solid which was recrystallised from ethyl acetate (yield = 87mg, 78%).

NMR (200MHz, DMSO-D6): δ 1.84(s,3H), 3.42(t,2H), 3.60(t,3H), 3.64(s,3H), 3.75(d of d,1H), 4.05(d,2H), 4.12(t,1H), 4.70(m,1H), 6.14(s,1H), 7.47(AB,4H), 8.24(m,1H).

5 MS: ESP+ (M+H) = 374.3.

Analysis: Calculated for $C_{19}H_{23}N_3O_5$: C, 61.1; H, 6.2; N, 11.1 %.

Found: C, 60.9; H, 5.9; N, 11.0 %.

Example 4 : N-((5S)-3-(4-(1-Methylsulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxo-
10 **oxazolidin-5-ylmethyl)acetamide**

Triethylamine (253mg), followed by a solution of methanesulfonyl chloride (69mg) in dichloromethane (1ml) was added, with ice-bath cooling, to a stirred suspension of N-((5S)-3-(4-(1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide (215mg, Example 2) in dichloromethane (10ml). TLC showed complete reaction after 2 hours. The
15 reaction mixture was washed with 2N. HCl, water, saturated sodium hydrogen carbonate and brine. The organic phase was dried over anhydrous sodium sulfate and evaporated to give the title compound as a crystalline solid which was recrystallised from acetonitrile (yield=117mg, 60%).

NMR (300MHz, DMSO-D6): δ 1.80(s, 3H), 2.55(broad, 2H), 2.90(s, 3H), 3.35(m, 4H),
20 3.73(d of d, 1H), 3.85(d, 2H), 4.10(t, 1H), 4.70(m, 1H), 6.15(s, 1H), 7.48(AB, 4H), 8.21(t, 1H).

MS: ESP+ (M+H)=394.

Analysis: Calculated for $C_{18}H_{23}N_3O_4S$: C, 54.9; H, 5.9; N, 10.7 %.

Found: C, 54.5; H, 5.6; N, 10.7 %.

25

Example 5 : N-((5S)-3-(4-(1-Acetyloxymethylcarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Sodium hydrogen carbonate (1.89g) was added to a stirred solution of Example 2 (1.92g) in a mixture of acetone (25ml) and water (12.5ml), and the mixture cooled to 0-3°C.

30 Acetoxyacetyl chloride (1.23g, 0.97ml) was added dropwise over 5-10 mins. and the reaction

mixture was stirred at 0-5°C for 30 minutes then allowed to warm to ambient temperature over 1 hour. TLC (10% methanol in dichloromethane) showed complete reaction.

The reaction mixture was diluted with water and extracted well with ethyl acetate. The organic phase was washed with brine and dried over anhydrous sodium sulfate. The title
5 compound crystallized during evaporation of the solvent and was triturated with ether (yield=1.70g, 91%).

NMR (300MHz, DMSO-D6): δ 1.81(s, 3H), 2.08(s, 3H), 2.5(obsured by DMSO), 3.40(t, 2H), 3.55(t, 1H), 3.65(t, 1H), 3.75(d of d, 1H), 4.10(m, 3H), 4.70(m, 1H), 4.85(d, 2H), 6.15(s, 1H), 7.48(m, 4H), 8.20(t, 1H).

10 MS: ESP+ (M+H)=416.

Example 6 : N-((5S)-3-(4-(1-Hydroxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Example 5 (1.70g) was stirred at ambient temperature with saturated methanolic ammonia
15 (20ml) for 18 hours. The initial suspension gave a solution after several hours from which the product crystallized. TLC showed complete reaction. The title compound was filtered off and washed with a little methanol (yield=1.28g, 84%).

NMR (300MHz, DMSO-D6): δ 1.81(s, 3H), 2.5(obsured by DMSO), 3.39(t, 2H), 3.50(m, 1H), 3.72(m, 2H), 4.10(m, 5H), 4.55(d, 1H), 4.70(m, 1H), 6.15(d, 1H),
20 7.45(m, 4H), 8.21(t, 1H).

MS: ESP+ (M+H)=374.

Analysis : Calculated for C₁₉H₂₃N₃O₅: C, 61.1; H, 6.2; N, 11.3 %.

Found: C, 60.8; H, 6.3; N, 11.1 %.

25 **Example 7 : N-((5S)-3-(3-Fluoro-4-(1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide**

Reference Example 2 (4.1g) was dissolved in TFA (10ml) and the yellow solution was heated briefly to reflux. The bulk of the TFA was evaporated and the residue was triturated with ether, precipitating initially a sticky gum which solidified to a yellow powder. The title
30 compound was obtained by filtration as a TFA salt, washed with ether and dried under vacuum (yield = 2.64g, 62%).

NMR (300MHz, DMSO-D6): δ 1.84(s, 3H), 2.65(s, 2H),
3.40(m, partially obscured by H₂O, 4H), 3.75(m, 3H), 4.10(t, 1H), 4.75(m, 1H), 6.05(m, 1H),
7.42(m, 3H), 8.25(t, 1H), 8.85(s, 2H).

5 **Example 8 : N-((5S)-3-(3-Fluoro-4-(1-acetyloxymethylcarbonyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide**

The title compound was prepared using a similar method to that describing the preparation of Example 5, except using Example 7 as the starting material (2.41g). Yield = 2.2g, 94%.

NMR (300MHz, DMSO-D6): δ 1.82(s, 3H), 2.10(s, 3H), 2.42(s, 2H), 3.40(m, 2H),
10 3.55(m, 1H), 3.62(m, 1H), 3.72(m, 1H), 4.10(m, 3H), 4.72(m, 1H), 4.82(d, 2H),
6.0(m, 1H), 7.38(m, 3H), 8.21(t, 1H).

MS: ESP+ (M+H)=434.

15 **Example 9 : N-((5S)-3-(3-Fluoro-4-(1-hydroxyacetyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide**

The title compound was prepared using a similar method to that describing the preparation of Example 6, except using Example 8 as the starting material (1.92g). Yield = 1.25g, 72%.

NMR (300MHz, DMSO-D6): δ 1.80(s, 3H), 2.45(observed by DMSO), 3.52(t, 1H),
3.65(t, 1H), 3.75(d of d, 1H), 4.10(m, 5H), 4.55(d, 1H), 4.70(m, 1H), 6.00(d, 1H),
20 7.24(d of d, 1H), 7.40(t, 1H), 7.47(d, 1H), 8.2(t, 1H).

MS: ESP+ (M+H)=392.

Example 10 : N-((5S)-3-(4-(2,3-Dihydro-6H-pyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

25 Triphenylarsine (61mg), Pd₂(dba)₃ (45.8mg) and lithium chloride (128mg) were added to a stirred solution of Reference Example 3 (278mg) in NMP (5ml) and the vessel was purged well with argon. After stirring for 5 minutes at ambient temperature, Reference Example 1 (397mg) was added and the reaction mixture was stirred at 40°C for 24 hours. TLC (ethyl acetate) indicated complete reaction. A 2.0M aqueous potassium fluoride solution (1ml) was
30 added and the reaction mixture was stirred at ambient temperature for 30 minutes. Water was added and the product was extracted with ethyl acetate, with filtration of the two phase

mixture. The organic phase was washed with water, brine, dried over anhydrous sodium sulfate and evaporated to a gum. The title compound was isolated by MPLC (Merck 9385 silica, using as eluant a mixture of acetonitrile (40%) and ethyl acetate), and was triturated with ether giving a crystalline powder (yield = 104mg, 33%).

- 5 NMR (300MHz, DMSO-D6): δ 1.83(t, 3H), 2.42(s, 2H), 3.40(m, 2H), 3.73(m, 1H), 3.80(t, 2H), 4.10(t, 1H), 4.20(d, 2H), 4.70(m, 1H), 6.22(s, 1H), 7.47(AB, 4H), 8.23(t, 1H).
MS: ESP+ (M+H)=317.

Example 11 : N-((5S)-3-(3-Fluoro-4-(2,3-dihydro-6H-pyran-4-yl)phenyl)-2-

10 **oxooxazolidin-5-ylmethyl)acetamide**

The title compound was prepared using a similar method to that describing the preparation of Example 10, except that N-((5S)-3-(3-fluoro-4-trimethyltinphenyl)-2-oxooxazolidin-5-ylmethyl)acetamide (415mg, WO Patent 94-13649, 23.06.94) was used as the starting material and a reaction time of 2 days was required. Yield = 130mg, 39%.

- 15 NMR (300MHz, DMSO-D6): δ 1.82(t, 3H), 2.40(s, 2H), 3.40(m, 2H), 3.71(d of d, 1H), 3.80(t, 2H), 4.10(t, 1H), 4.20(d, 2H); 4.73(m, 1H), 6.08(s, 1H), 7.28(d of d, 1H), 7.38(t, 1H), 7.47(d of d, 1H), 8.21(t, 1H).
MS: ESP+ (M+H)=335.

20 **Example 12 : N-((5S)-3-(3-Fluoro-4-(1-benzyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide**

Reference Example 8 (6.9g) was dissolved in TFA (10ml) and heated briefly to reflux. The TFA was evaporated and the residue was basified with aqueous ammonium hydroxide solution and the product extracted with ethyl acetate. The organic phase was dried over

- 25 anhydrous magnesium sulfate and evaporated to an oil which crystallized on trituration with ether to give the title compound (yield = 3.78g, 75% over 2 stages).

NMR (300MHz, DMSO-D6): δ 1.81(s, 3H), 2.41(s, 2H), 2.60(m, 2H), 3.05(s, 2H), 3.40(m, 2H), 3.58(s, 2H), 3.72(m, 1H), 4.10(m, 1H), 4.70(m, 1H), 5.95(s, 1H), 7.31(m, 10H), 8.20(t, 1H).

- 30 MS: ESP+ (M+H)=424.

Example 13 : (5R)-Hydroxymethyl-3-(3-fluoro-4-(N-benzyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)oxazolidin-2-one

n-Butyllithium/hexane (28.1ml of 1.40M) was slowly added to a stirred partial solution of Reference Example 7 (8.7g) in dry THF (100ml) at -70°C under argon. The solution was stirred for 15 minutes, then a solution of R-glycidyl butyrate (3.04g) in THF (15ml) was added and the reaction mixture was allowed to warm to ambient temperature and stirred for 18 hours. The reaction mixture was quenched with water and extracted with ethyl acetate. The organic phase was washed with water, brine, dried over anhydrous sodium sulfate and evaporated to give the title compound as a gum which was isolated by MPLC (silica, using as
10 eluant a mixture of methanol (5%) and dichloromethane) and recrystallised from acetonitrile. Yield = 4.78g, 58%.

NMR (300MHz, DMSO-D6): δ 2.43(s, 2H), 2.60(m, 2H), 3.05(s, 2H), 3.53(m, 1H), 3.58(s, 2H), 3.65(m, 1H), 3.80(d of d, 1H), 4.07(t, 1H), 4.70(m, 1H), 5.19(t, 1H), 5.95(s, 1H), 7.30(m, 7H), 7.47(d of d, 1H).

15 MS: ESP+ (M+H)=383.

Analysis: Calculated for $C_{22}H_{23}N_2O_3F$: C, 69.0; H, 6.0; N, 7.3 %.

Found: C, 69.2; H, 6.1; N, 7.4 %.

Example 14 : N-((5S)-3-(3-Fluoro-4-(1-methoxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

A solution of the hydrochloride salt of Example 7 (300 mg) and $NaHCO_3$ (340mg) in acetone (8ml) / water (4ml) was cooled to 0-4°C and methoxyacetyl chloride (176mg) was added dropwise. After stirring for 15 minutes additional $NaHCO_3$ and methoxyacetyl chloride (as above) were added giving complete reaction within 10 minutes. The reaction mixture was
25 diluted with water and extracted with ethyl acetate. The organic phase was washed with saturated NaCl solution, dried over anhydrous Na_2SO_4 and evaporated to a gum. The title compound was isolated by MPLC (4% MeOH / CH_2Cl_2 , Merck 9385 silica), and was triturated with ether (yield = 236mg, 72%).

NMR (300MHz, DMSO-D6) δ : 1.80(s, 3H), 2.4(broad, 2H), 3.27(s, 3H+ water),
30 3.38(t, 2H), 3.52(broad, 1H), 3.62(broad, 1H), 3.71(q, 1H), 4.18(m, 5H), 4.70(m, 1H), 5.97(s, 1H), 7.25(d of d, 1H), 7.35(t, 1H), 7.45(d of d, 1H), 8.19(t, 1H).

MS: ESP+ (M+H) = 406.

The hydrochloride salt of Example 7 was prepared by the reaction in dry dichloromethane at 0-4°C of Example 12 (16.6g) with 1-chloroethyl chloroformate (6.73g, added dropwise), in the presence of triethylamine (0.59g). After reaction for 0.5 hours the solvent was removed under vacuum and the residue refluxed with methanol (100ml) for 0.5 hours. The methanol was removed under vacuum and the residue recrystallised from ethanol (250ml) to yield the hydrochloride salt of Example 7 (11.92g, 82%, NMR as previously assigned for Example 7).

10 **Example 15 : N-((5S)-3-(3-Fluoro-4-(1-methyloxalyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide**

The hydrochloride salt of Example 7 (300mg) was suspended in ethyl acetate, washed with aqueous ammonia, dried over anhydrous Na₂SO₄ and evaporated to give the free base of Example 7. Triethylamine (152mg) was added to a mixture of the resulting free base of Example 7 in dichloromethane (10ml), followed by the addition of methyloxalyl chloride (123mg). Reaction was complete after 10 minutes. The reaction mixture was washed with 2N HCl, water, aqueous ammonia and saturated NaCl. The organic phase was dried over anhydrous Na₂SO₄ and evaporated to a gum. The title compound was isolated by MPLC (4% MeOH/CH₂Cl₂, Merck 9385 silica), and was triturated with ether (yield = 181mg, 53%).

15 NMR (300MHz, DMSO-D₆) δ : 1.79(s, 1H), 2.50(obsured by solvent), 3.38(t, 2H), 3.55(t, 1H), 3.70(m, 2H), 3.82(d, 3H), 4.05(s, 1H), 4.12(m, 2H), 4.70(m, 1H), 5.98(d, 1H), 7.25(d of d, 1H), 7.35(t, 1H), 7.45(d of d, 1H), 8.19(t, 1H).

MS: ESP+ (M+H) = 420.

25 **Example 16: N-((5S)-3-(3-Fluoro-4-(ethylaminocarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide**

Triethylamine (101mg), followed by ethyl isocyanate (71mg) were added to a stirred suspension of the hydrochloride salt of Example 7 (300 mg) in THF (10ml). After stirring for 2 hours, additional triethylamine and ethyl isocyanate (as above) were added, giving complete reaction after 3 days at ambient temperature. The reaction mixture was washed with 2N HCl, water, saturated NaHCO₃ solution and saturated NaCl solution. Crystallised product was

filtered off. The organic phase was dried over anhydrous Na_2SO_4 and evaporated to give a crystalline solid. The combined product was purified by MPLC (using a mixture of methanol and dichloromethane of increasing polarity as eluant, Merck 9385 silica), and was triturated with ether to give the title compound (222mg, 68%).

5 NMR (300MHz, DMSO-D6) δ : 1.00(t, 3H), 1.80(s, 3H), 2.37(broad, 2H), 3.05(6 line, 2H), 3.38(t, 2H), 3.47(t, 2H), 3.71(q, 1H), 3.92(broad, 2H), 4.10(t, 1H), 4.70(t, 1H), 5.98(s, 1H), 6.72(m, 1H), 7.25(d of d, 1H), 7.35(t, 1H), 7.45(d of d, 1H), 8.20(t, 1H).

MS: ESP+ (M+H) = 405.

10 Example 17: N-((5S)-3-(3-Fluoro-4-(1-acetoacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Triethylamine (202mg), followed by diketene (227mg) were added to a stirred suspension of the hydrochloride salt of Example 7 (665mg) in dichloromethane (15ml). The starting material dissolved and a precipitate of product began to separate after 15 minutes. The reaction was complete after 30 minutes. The precipitate was filtered off, washed with dichloromethane and recrystallised from acetonitrile to give the title compound (520mg, 69%).

NMR (300MHz, DMSO-D6) δ : 1.81(s, 3H), 2.15(d, 3H), 2.40(broad, 2H), 3.39(t, 2H), 3.50(t, 1H), 3.70(m, 4H), 4.10(m, 3H), 4.72(m, 1H), 5.98(d, 1H), 7.25(d of d, 1H), 7.38(t of d, 1H), 7.45(d, 1H), 8.20(t, 1H).

MS: ESP+ (M+H) = 418.

Example 18 : N-((5S)-3-(3-Fluoro-4-(1-(2-{2-methoxyethoxy}ethoxy)methylcarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

25 N-Methylmorpholine (101mg), 2-[2-(2-methoxyethoxy)ethoxy]acetic acid (154mg) and N-hydroxybenzotriazole (122mg, allowing for 5% water) were added to a mixture of the hydrochloride salt of Example 7 (277mg) and DMF (5ml). The mixture was stirred for 5 minutes at ambient temperature and then 1-ethyl-3-dimethylaminopropyl-carbodiimide (165mg) was added and the mixture stirred for 18 hours. The solvent was evaporated under high vacuum and the residue partially purified by MPLC [using a mixture of methanol and dichloromethane increasing in polarity from 5% to 25% methanol as eluant, Merck 9385

silica]. Final purification by reverse-phase MPLC [using a mixture of acetonitrile and water, increasing in acetonitrile from 0% to 25% as eluant, HP20SS resin] and trituration with ether gave the title compound (111mg, 30%).

NMR (300 MHz, DMSO-D₆) δ : 1.80(s, 1H), 1.86(s, 1H), 2.43(partially obscured), 3.20(s, 2H), 3.40(m, 3H), 3.55(m, 10H), 3.72(m, 1H), 4.08(m, 3H), 4.20(s, 2H), 4.70(m, 1H), 6.00(s, 1H), 7.26(d, 1H), 7.35(t, 1H), 7.45(d, 1H), 8.20(t, 1H).

MS: ESP+ (M+H) = 494.

Example 19: N-((5S)-3-(3-Fluoro-4-(1-formyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

In an analogous manner to that described in Example 15, the hydrochloride salt of Example 7 (370mg) was converted to the free base, which was then refluxed with ethyl formate (25ml) for 24hours. The reaction mixture was evaporated and the title compound was isolated by MPLC (5% MeOH / CH₂Cl₂ eluant, Merck 9385 silica). The title product crystallised on trituration with ether (297mg, 82%).

NMR (300MHz, DMSO-D₆) δ : 1.80(s, 1H), 2.43(m, 2H), 3.40(t, 2H), 3.58(m, 2H), 3.72(q, 1H), 4.02(d, 1H), 4.07(s, 1H), 4.11(t, 1H), 4.72(m, 1H), 5.98(d, 1H), 7.25(d of d, 1H), 7.35(t, 1H), 7.45(d of d, 1H), 8.12(d, 1H), 8.20(t, 1H).

MS: ESP+ (M+H) = 362.

20

Example 20: N-((5S)-3-(3-Fluoro-4-(1-acetylaminoacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

N-Methylmorpholine (101mg), N-acetyl glycine (101mg) and N-hydroxybenzotriazole (122mg, allowing for 5% water) were added to a mixture of the hydrochloride salt of Example 7 (277mg) and DMF (5ml). The mixture was stirred for 5 minutes at ambient temperature then 1-ethyl-3-dimethylaminopropyl-carbodiimide (165mg) was added and the mixture was stirred for 18 hours. The solvent was evaporated under high vacuum and the title compound was isolated by MPLC (using a mixture of methanol and dichloromethane increasing in polarity from 4-10% methanol as eluant, Merck 9385 silica). The title product crystallised on trituration with ether (270mg, 83%).

30

NMR (300MHz, DMSO-D6) δ : 1.80(s, 1H), 1.86(s, 1H), 2.45(partially obscured), 3.39(t, 2H), 3.60(m, 2H), 3.72(q, 1H), 3.95(d of d, 2H), 4.10(m, 3H), 4.72(m, 1H), 5.99(broad, 1H), 7.25(d of d, 1H), 7.38(t of d, 1H), 7.45(d, 1H), 7.96(broad, 1H), 8.20(t, 1H).

MS: ESP+ (M+H) = 433.

5

Example 21: N-((5S)-3-(3-Fluoro-4-(-acetylacetoxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

4-Dimethylaminopyridine (4mg) followed by diketene (118mg) were added to a stirred solution of the Example 9 (261mg) in dry dichloromethane (10ml). After stirring for 30
10 minutes at ambient temperature the reaction mixture was evaporated and the title compound was isolated by MPLC (5% MeOH / CH₂Cl₂ eluant, Merck 9385 silica). The title product crystallised on trituration with ether (255mg, 80%).

NMR (300MHz, DMSO-D6) δ : 1.80(s, 3H), 2.22(s, 3H), 2.50(obsured), 3.40(t, 2H), 3.55(t, 1H), 3.64(s, 3H), 3.73(q, 1H), 4.07(m, 3H), 4.68(m, 1H), 4.87(d, 2H), 6.13(broad, 1H),
15 7.46(m, 4H), 8.20(t, 1H).

MS: ESP+ (M+H) = 458.

Example 22 : N-((5S)-3-(3-Fluoro-4-(1-(2-methyl-2-phenylamino)ethenylcarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

20 A mixture of Example 17 (104mg), aniline (47mg), p-toluenesulfonic acid (p-TSA) (5mg) and ethanol (ca. 10ml) was stirred under reflux for 24 hours. Additional aniline (233mg) and p-TSA (5mg) were added, together with 3A molecular sieve powder (1g) and reflux continued for a further 24 hours. The reaction mixture was filtered, evaporated. MPLC [using a mixture of acetonitrile and ethyl acetate, increasing in acetonitrile from 25% to 50% as eluant], and
25 trituration with ether gave the title compound (43mg, 35%).

NMR (300 MHz, DMSO-D6) δ : 1.82(s, 3H), 2.05(s, 3H), 2.45(obsured), 3.40(q, 2H), 3.70(m, 3H), 4.14(m, 3H), 4.72(m, 1H), 5.08(s, 1H), 6.04(s, 1H), 7.10(m, 3H), 7.37(m, 5H), 8.20(t, 1H), 11.67(s, 1H).

MS: ESP+ (M+H) = 493.

30

Example 23 : There is no compound with Example No. 23.

Example 24 : N-((5S)-3-(3-fluoro-4-(1-cyano-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

4-Dimethylaminopyridine (24mg), triethylamine (606mg) and a solution of cyanogen bromide (1.06g) in dichloromethane (5ml) were added to a stirred suspension of the hydrochloride salt of Example 7 (739mg) in dry dichloromethane (15ml). After stirring for 1 hour at ambient temperature, a solution had formed and the reaction was complete. The reaction mixture was washed with 2N HCl, aqueous NaHCO₃ and dried over anhydrous Na₂SO₄. The mixture was evaporated to give a gum which crystallised on trituration with ether (488mg, 68%).

NMR (300MHz, DMSO-D6) δ : 1.81(s, 3H), 2.50(d, obscured), 3.39(t, 4H), 3.71(q, 1H),
10 3.92(d, 2H), 4.11(t, 1H), 4.72(m, 1H), 5.95(s, 1H), 7.27(d of d, 1H), 7.35(t, 1H), 7.47(d of d, 1H), 8.20(t, 1H).

MS: ESP+ (M+H) = 359.

Example 25: N-((5S)-3-(3-Fluoro-4-(1-(pyrimid-2-yl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

A stirred mixture of the hydrochloride salt of Example 7 (1.29g), NaHCO₃ (1.18g) and 2-chloropyrimidine (601mg) was heated under reflux for 5 hours. The solvent was evaporated and the residue was taken into ethyl acetate and washed with water and brine. The organic phase was dried over anhydrous Na₂SO₄ and evaporated. The title compound was isolated by
20 MPLC (using a mixture of acetonitrile and ethyl acetate increasing in polarity from 50% to 65% acetonitrile as eluant, Merck 9385 silica), and was triturated with ether (939mg, 65%).

NMR (300MHz, DMSO-D6) δ : 1.82(s, 3H), 2.50(obscured), 3.40(t, 2H), 3.70(q, 1H), 3.95(t, 2H), 4.10(t, 1H), 4.30(q, 2H), 4.70(m, 1H), 6.10(s, 1H), 6.62(t, 1H), 7.27(d of d, 1H), 7.39(t, 1H), 7.47(d of d, 1H), 8.20(t, 1H), 8.37(d, 2H).

25 MS: ESP+ (M+H) = 412.

Example 26: N-((5S)-3-(3-Fluoro-4-(1-(tetrazol-5-yl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Azidotrimethylsilane (154mg) and dibutyltin oxide (97mg) were added to a stirred suspension
30 of Example 24 (240mg) in dry toluene (10ml), and the mixture stirred at 70°C for 18 hours.

On cooling, methanol (5ml) was added and after stirring for 10 minutes the solvent was

evaporated. The title compound was isolated by MPLC (20% MeOH / CH₂Cl₂, Merck 9385 silica) and crystallised on evaporating the fractions to a small volume. The title product was filtered off washed with a little cold methanol (150mg, 56%).

NMR (300MHz, DMSO-D6) δ : 1.80(s, 3H), 2.55(s, 2H), 3.30(obsured, 1H), 3.39(t, 2H),
5 3.63(t, 2H), 3.73(q, 1H), 4.05(d, 2H), 4.12(t, 1H), 4.62(m, 1H), 6.08(s, 1H), 7.27(d of d, 1H),
7.39(t, 1H), 7.47(d of d, 1H), 8.19(t, 1H).

MS: ESP+ (M+H) =402.

Example 27 : There is no compound with Example No. 27.

10

Example 28 : There is no compound with Example No. 28.

Example 29: N-((5S)-3-(3-Fluoro-4-(1-methylsulfonyl-1,2,5,6-tetrahydro-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

15 Triethylamine (242mg), followed by methanesulfonyl chloride (115mg) were added to a stirred suspension of the hydrochloride salt of Example 7 (296mg) in dichloromethane (10ml). After stirring for 30 minutes, additional triethylamine and methanesulfonyl chloride (as above) were added giving a solution, and complete reaction after a further 30 minutes at ambient temperature. The reaction mixture was washed with 2N HCl, water, saturated
20 NaHCO₃ and saturated NaCl. The organic phase was dried over anhydrous Na₂SO₄ and evaporated to give a crystalline solid. This was purified by MPLC (using 5% MeOH / CH₂Cl₂ as eluant, Merck 9385 silica) and triturated with ether to give the title compound (209mg, 64%).

NMR (300MHz, DMSO-D6) δ : 1.80(s, 3H), 2.55(s, 2H), 2.93(s, 3H), 3.35(m, 4H),
25 3.72(q, 1H), 3.85(d, 2H), 4.11(t, 1H), 4.72(m, 1H), 6.02(s, 1H), 7.27(d of d, 1H),
7.39(t, 1H), 7.47(d of d, 1H), 8.20(t, 1H).

MS: ESP+ (M+H) = 412.

Example 30 : N-((5S)-3-(4-(1-Decyloxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Sodium hydride (29mg, 60% in mineral oil) was added to a stirred solution of Example 6 (242mg) in dry DMF (5ml). There was an effervescence and a precipitate formed. After 5 minutes, decyl iodide (209mg) was added and the reaction mixture was stirred for 24 hours at ambient temperature. Additional sodium hydride and decyl iodide (as above) were added giving almost complete reaction after a further 24 hours. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic phase was washed with water and saturated NaCl, was dried over anhydrous Na₂SO₄ and evaporated to give a gum. The title compound was isolated by MPLC (using 3.5% MeOH / CH₂Cl₂ as eluant, Merck 9385 silica), and was triturated with ether. Yield = 132mg, 40%.

MS: ESP+ (M+H) = 514.

The title compound was characterised by an HPLC retention time of 23.30 mins., using the HPLC conditions described in Example 35.

15

Example 31 : N-((5S)-3-(4-(1-Butyryloxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Triethylamine (101mg) was added to a stirred solution of Example 6 (242mg) in dichloromethane (10ml), followed by the dropwise addition of butyryl chloride (76mg) in dichloromethane (0.2ml). After stirring for 30 minutes, additional triethylamine and butyryl chloride (as above) were added giving complete reaction after a further 30 minutes at ambient temperature. The reaction mixture was washed with 2N HCl, water, saturated NaHCO₃, and saturated NaCl. The organic phase was dried over anhydrous Na₂SO₄ and evaporated to a crystalline solid. This was purified by MPLC (using 4% MeOH / CH₂Cl₂ as eluant, Merck 9385 silica) and triturated with ether to give the title compound. Yield = 237mg, 82%.

25

NMR (300MHz, DMSO-D6) δ : 0.90(t, 3H), 1.55(m, 2H), 1.81(s, 3H), 2.32(t, 2H), 2.55(obsured), 3.40(t, 2H), 3.56(t, 1H), 3.64(t, 1H), 3.75(q, 1H), 4.10(m, 3H), 4.69(m, 1H), 4.83(d, 2H), 6.15(broad, 1H), 7.48(m, 4H), 8.20(t, 1H).

MS: ESP+ (M+H) = 443.

30

Example 32 : N-((5S)-3-(4-(1-Decanoyloxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Using an analogous procedure to that described in Example 31, Example 6 was acylated with decanoyl chloride (2 x 149mg) to give the title compound. Yield = 268mg, 78%.

5 NMR (300MHz, DMSO-D6) δ : 0.90(t, 3H), 1.23(s, 12H), 1.55(m, 2H), 1.81(s, 3H), 2.32(t, 2H), 2.55(obsured), 3.40(t, 2H), 3.56(t, 1H), 3.64(t, 1H), 3.75(q, 1H), 4.10(m, 3H), 4.69(m, 1H), 4.83(d, 2H), 6.15(broad, 1H), 7.48(m, 4H), 8.20(t, 1H).

MS: ESP+ (M+H) = 528.

10 **Example 33 : N-((5S)-3-(4-(1-Ethylaminocarboxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide**

Triethylamine (101mg), followed by ethyl isocyanate (142mg) were added to a stirred solution of Example 6 (298 mg) in dichloromethane (10ml). After stirring for 18 hours at ambient temperature, additional triethylamine and ethyl isocyanate (as above) were added
15 giving complete reaction after a further 24 hours. A precipitate of product was filtered off and a second crop was obtained on reducing the volume. The combined material was recrystallised from ethanol to give the title compound. Yield = 251mg, 71%.

NMR (300MHz, DMSO-D6) δ : 1.00(t, 3H), 1.80(s, 3H), 2.50(obsured), 3.00(m, 2H), 3.40(t, 2H), 3.55(t, 1H), 3.65(t, 1H), 3.73(q, 1H), 4.07(m, 3H), 4.68(m, 3H), 6.13(broad, 1H),
20 7.27(broad, 1H), 7.50(m, 4H), 8.21(t, 1H).

MS: ESP+ (M+H) = 445.

Example 34: N-((5S)-3-(4-(1-Benzyloxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

25 Using an analogous procedure to that described in Example 31, Example 6 was acylated on a 2.0mM scale with benzyloxyacetyl chloride (3 x 461mg) to give the title compound. Yield = 705mg, 68%. NMR (300MHz, DMSO-D6) δ : 1.82(s, 3H), 2.50(obsured), 3.40(t, 2H), 3.58(t, 1H), 3.66(t, 1H), 3.73(q, 1H), 4.10(m, 3H), 4.24(s, 2H), 4.58(s, 2H), 4.70(m, 1H), 4.95(d, 2H), 6.15(broad, 1H), 7.35(m, 5H), 7.50(m, 4H), 8.20(t, 1H).

30 MS: ESP+ (M+H) = 522.

Example 35: N-((5S)-3-(4-(1-(4-*t*-Butylcyclohexylcarbonyl))-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Using an analogous procedure to that described in Example 20, the TFA salt of Example 2 was reacted with 4-tert-butylcyclohexanecarboxylic acid in place of N-acetylglycine, to give the title compound. Yield = 208mg, 58%.

MS: ESP+ (M+H) = 482.

The title compound was characterised by an HPLC retention time of 21.72 mins., using the following HPLC conditions : Column: Hypersil ODS 5m; Flow rate: 1.5 ml/min.; UV: 254nm; gradient parameters: Solvent A 1 mM TFA / water, Solvent B 1 mM TFA / acetonitrile; time 0 95% A / 5% B; time 3 mins. 95% A / 5% B; time 17 mins. 5% A / 95% B; time 19 mins 95% A / 5% B; time 20 mins 95% A / 5% B.

Example 36: N-((5S)-3-(4-(1-(2-Hydroxy-decanoyl))-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Using an analogous procedure to that described in Example 20, the TFA salt of Example 2 was reacted with 2-hydroxydecanoic acid in place of N-acetylglycine, to give the title compound. Yield = 252mg, 65%.

MS: ESP+ (M+H) = 514.

The title compound was characterised by an HPLC retention time of 22.80 mins., using the HPLC conditions described in Example 35.

Example 37: N-((5S)-3-(4-(1-(Pyrimid-2-yl))-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Using an analogous procedure to that described in Example 25, the TFA salt of Example 2 was reacted on a 1.5mM scale with 2-chloropyrimidine, to give the title compound. Yield = 220mg, 36%.

NMR (300MHz, DMSO-D₆) δ : 1.80(s, 3H), 2.50(obsured), 3.40(t, 2H), 3.73(q, 1H), 3.98(t, 2H), 4.10(t, 1H), 4.30(d, 2H), 4.69(m, 1H), 6.25(s, 1H), 6.62(t, 1H), 7.48(m, 4H), 8.20(t, 1H), 8.37(d, 2H).

MS: ESP+ (M+H) = 394.

Example 38: N-((5S)-3-(4-(1-p-Toluenesulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Triethylamine (202mg), followed by p-toluenesulfonyl chloride (171mg) were added to a stirred suspension of the TFA salt of Example 2 (322mg) in dichloromethane (10ml). After stirring for 30 minutes at ambient temperature, the reaction mixture had become a solution and the reaction was complete. The solution was washed with 2N HCl, water, saturated NaHCO₃, and saturated NaCl. The organic phase was dried over anhydrous Na₂SO₄ and evaporated to give a crystalline solid which was triturated with ether to give the title compound. Yield = 306mg, 87%.

10 NMR (300MHz, DMSO-D6) δ : 1.79(s, 3H), 2.35(s, 3H), 2.50(obsured), 3.17(m, 2H), 3.38(m, 2H), 3.64(broad, 2H), 3.70(m, 1H), 4.08(m, 1H), 4.68(m, 1H), 6.05(broad, 1H), 7.40(m, 6H), 7.68(m, 2H), 8.20(t, 1H).

MS: ESP+ (M+H) =470.

15 **Example 39: N-((5S)-3-(4-(1-Hydroxyacetyl-1,2,3,4,5,6-hexahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide**

A mixture of Example 6 (242mg) and ethanol (10ml) was hydrogenated over 10% palladium on charcoal (25mg) at atmospheric pressure for 3 hours. The catalyst was removed by Millipore filtration and the solvent was evaporated leaving an oil which crystallised on trituration with ether to give the title compound. Yield = 230mg, 95%.

NMR (300MHz, DMSO-D6) δ : 1.50(m, 2H), 1.75(m, 2H), 1.80(s, 3H), 2.70(m, 2H), 3.04(t, 1H), 3.37(t, 2H), 3.73(m, 2H), 4.09(m, 3H), 4.25(m, 2H), 4.69(m, 1H), 7.23(d, 2H), 7.42(d, 2H), 8.20(t, 1H).

MS: ESP+ (M+H) =376.

25

Example 40: N-((5S)-3-(3-Fluoro-(2,3,4,5-tetrahydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Using an analogous procedure to that described in Example 39, Example 11 was hydrogenated on a 0.6mM scale to give the title compound in quantitative yield.

NMR (300MHz, DMSO-D6) δ : 1.65(m, 4H), 1.80(s, 3H), 3.00(m, 1H), 3.36(t, 2H), 3.44(d of d, 2H), 3.70(d of d, 1H), 3.90(d of d, 2H), 4.07(t, 1H), 4.70(m, 1H), 7.23(d of d, 1H), 7.32(t, 1H), 7.41(d of d, 1H), 8.20(t, 1H).

MS: ESP+ (M+H) =337.

5

Example 41: N-((5S)-3-(3,5-Difluoro-(2,3,4,5-tetrahydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Using an analogous procedure to that described in Example 39, Example 47 was hydrogenated on a 4.23mM scale to give the title compound (1.38g, 93%).

10 NMR (300MHz, DMSO-D6) δ : 1.55(d of d, 2H), 1.95(m, 2H), 3.13(m, 1H), 3.38(m, 4H), 3.69(q, 1H), 3.91(d of d, 2H), 4.08(t, 1H), 4.72(m, 1H), 7.25(d, 2H), 8.18(t, 1H).

MS: ESP+ (M+H) =355.

Example 42: N-((5S)-3-(3-Fluoro-4-(1-hydroxyacetyl-1,2,3,4,5,6-hexahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

Using an analogous procedure to that described in Example 39, Example 9 was hydrogenated on the scale of Example 39 to give the title compound (236mg, 92%).

NMR (300MHz, DMSO-D6) δ : 1.60(m, 4H), 1.80(s, 3H), 2.69(t, 1H), 3.04(m, 2H), 3.39(t, 2H), 3.73(m, 2H), 4.08(m, 3H), 4.48(m, 2H), 4.70(m, 1H), 7.20(d of d, 1H),
20 7.30(t, 1H), 7.43(d of d, 1H), 8.19(t, 1H).

MS: ESP+ (M+H) =394.

Example 43: N-((5S)-3-(4-(1-Hydroxyacetyloxyacetyl-1,2,3,4,5,6-hexahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide

25 A mixture of Example 34 (261mg) and ethanol (10ml) was hydrogenated over 10% palladium on charcoal (125mg) at atmospheric pressure for 3 days. The catalyst was removed by Millipore filtration and the solvent was evaporated leaving an oil which crystallised on trituration with ether to give the title compound. Yield = 179mg, 83%.

NMR (300MHz, DMSO-D6) δ : 1.55(m, 4H), 1.82(s, 3H), 2.70(m, 2H), 3.10(m, 1H),
30 3.39(t, 2H), 3.70(q, 1H), 3.80(m, 1H), 4.09(m, 3H), 4.41(m, 1H), 4.69(m, 1H), 4.84(s, 2H), 5.40(t, 1H), 7.24(d, 2H), 7.44(d, 2H), 8.20(t, 1H).

MS: ESP+ (M+H) = 434.

Example 44: N-((5S)-3-(3,5-Difluoro-4-(1,2,5,6-tetrahydro-4H-pyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

- 5 N-BOC-acetamide (3.1g, Chem. Pharm. Bull., 36, 3125 (1988)) was added to a solution of Reference Example 11 (5.3g) in THF (100ml), and the mixture cooled in an ice-bath. Tributyl phosphine (4.8ml, 3.89g) and 1,1'-(azodicarbonyl)-dipiperidine (4.9g) were added and the mixture allowed to stir at ice-bath temperature for 30 minutes after which a precipitate was evident. The reaction was then stirred overnight at ambient temperature. After filtration and
- 10 evaporation in vacuo the residue was purified on silica gel using a 1:1 mixture of isohexane and ethyl acetate as eluant to give a solid (6.77g). The solid was dissolved in TFA (10ml) and refluxed for a few minutes. The TFA was removed in vacuo and the residue dissolved in a mixture of ethyl acetate and methanol. The solvent was evaporated and trituration with ether gave the title product (3.5g, 58%) as the TFA salt (58%). The filtrate gave a further impure
- 15 gum (1.25g) which was suitable for further conversions.

NMR (300 MHz, DMSO-D6) δ : 2.50(broad s, 2H), 3.40(t, 2H), 3.71(q, 1H), 3.77(broad s, 2H), 4.11(t, 1H), 4.74(m, 1H), 5.89(broad s, 1H), 7.34(d, 1H), 8.20(t, 1H), 8.94(broad s, 2H).

20 **Example 45: N-((5S)-3-(3,5-Difluoro-4-(1-acetoxymethylcarbonyl-1,2,5,6-tetrahydro-4H-pyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.**

Using an analogous reaction and work-up procedure to that described in Example 5, the TFA salt of Example 44 and acetoxy-acetylchloride were reacted on a 7.16mM scale to give the title compound. Yield = 2.62g, 81%.

- 25 NMR (300MHz, DMSO-D6) δ : 1.81(s, 3H), 2.08(s, 3H), 2.30(broad s, 1H), 2.40(broad s, 1H), 3.40(t, 2H), 3.55(t, 1H), 3.64(t, 1H), 3.72(q, 1H), 4.08(m, 3H), 4.74(m, 1H), 4.83(d, 2H), 5.85(m, 1H), 7.30(d, 2H), 8.20(t, 1H).

MS: ESP+ (M+H) = 452.

Example 46: N-((5S)-3-(3,5-Difluoro-4-(1-hydroxyacetyl-1,2,5,6-tetrahydro-4-pyridyl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

Using an analogous procedure to that described in Example 6, Example 45 (0.5g) was converted to the title compound (solid 0.38g, 84%).

5 NMR (300MHz, DMSO-D6) d : 1.81(s, 3H), 2.33(broad d, 2H), 3.38(t, 2H), 3.51(t, 1H), 3.66(t, 1H), 3.70(q, 1H), 4.08(m, 3H), 4.12(t, 2H), 4.58(broad d, 1H), 4.74(m, 1H), 5.85(d, 1H), 7.30(d, 2H), 8.20(t, 1H).

MS: ESP+ (M+H) = 410.

10 **Example 47 : N-((5S)-3-(3,5-Difluoro-4-(2,3-dihydro-6H-pyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.**

N-BOC-acetamide (5.37g, Chem. Pharm. Bull., 36, 3125 (1988)), followed by tributyl phosphine (8.3ml) and 1,1'-(azodicarbonyl)-dipiperidine (8.52g) were added to a stirred mixture of Reference Example 14 (7.0g) and anhydrous THF (200ml) in an ice-bath. The
15 mixture was stirred for a further 30 minutes at 0-5°C and then stirred to ambient temperature over-night. The mixture was filtered, the solid washed with THF and the combined filtrates evaporated in vacuo and purified on silica gel (using a 4:6 mixture of isohexane and ethyl acetate as eluant). There was thus obtained an oil, which was dissolved in TFA (20ml) by slow addition of TFA to the oil, refluxed briefly for few minutes and evaporated in vacuo. The
20 residue was taken up in ethyl acetate (200ml) washed with saturated NaHCO₃, the extracts counter-washed and the combined organic extracts evaporated in vacuo. The residue was purified on silica gel (using a mixture of 5% methanol and dichloromethane as eluant) to give the title compound as a solid (5.4g, 68%).

NMR (300MHz, DMSO-D6) d : 1.82(s, 3H), 2.26(broad s, 2H), 3.18(t, 2H), 3.70(q, 1H),
25 3.78(t, 2H), 4.10(t, 1H), 4.18(q, 2H), 4.65(m, 1H), 5.89(s, 1H), 7.26(s, 1H), 7.33(s, 1H), 8.19(t, 1H).

MS: ESP+ (M+H) = 353.

Example 48: N-((5S)-3-(4-(2-Methyl-3,6- and -5,6-dihdropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

Reference Example 15 (0.49 g), palladium(0) bis(dibenzylideneacetone) (0.86 g), triphenylarsine (0.115 g) and lithium chloride (0.24 g) were dissolved in degassed *N*-methyl morpholine (33 ml). The stannane (Reference Example 1) (0.75 g) was added and the solution stirred at 40°C for 18 hours. The mixture was then treated with aqueous potassium fluoride (5 ml, 2M), stirred for 0.5 hours and filtered through Celite. The solution was drowned with water (250 ml) and extracted with ethyl acetate (2 x 30 ml). The combined organics were washed with water (2 x 100 ml), dried and evaporated to the crude product (0.333 g, 54%) which was purified by firstly silica-gel MPLC [using a mixture of 2% MeOH and CH₂Cl₂ as eluant] and then preparative TLC [using a mixture of 13% MeOH and CH₂Cl₂ as eluant] to give the title mixture of 2-methyl regioisomers (0.054g, 9%).

NMR (300 MHz, CDCl₃) d: 1.30-1.39 (m, 3H), 2.00-2.05 (s/s, 3H), 2.27-2.39 (broad m, 1.5H), 2.57-2.69 (m, 0.5H), 3.57-3.97 (4H, m), 4.00-4.20 (m, 2H), 4.30-4.40 (m, 1H), 4.72-4.81 (m, 1H), 6.00 (s, 0.5H), 6.05-6.10 (m, 0.5H), 6.40 (broad s, 1H), 7.39 (dd, 2H) and 7.45 (d, 2H).

MS: ESP+ (M+H) = 331.

Example 49: N-((5S)-3-(4-(2,6-Dimethyl-2,3-dihydro-6H-pyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

Reference Example 18 (0.656 g), palladium(0) bis(dibenzylideneacetone) (0.123 g), triphenylarsine (0.164 g) and lithium chloride (0.338 g) were dissolved in degassed *N,N*-dimethylformamide (18 ml). The stannane (Reference Example 1) (1.07 g) was added after 5 minutes and stirring was continued for 19 hours at 40 °C. Aqueous potassium fluoride (6.3 ml, 2M) was added and stirring continued for 15 minutes whereupon the mixture was filtered through Celite and evaporated to an oil. The oil was purified by silica-gel MPLC [using a mixture of ethyl acetate and isohexane increasing in polarity from 75% to 100% ethyl acetate as eluant] to afford a foam which was triturated with diethyl ether to give the title compound as a powder (0.679g, 78%).

NMR (300 MHz, CDCl₃) d : 1.33 and 1.35 (d/d, each 3H), 2.01 (s, 3H), 2.28-2.35 (m, 2H), 3.55-3.82 (m, 4H), 4.05 (t, 1H), 4.36-4.25 (m, 1H), 4.72-4.82 (m, 1H), 6.00 (d, 1H), 6.01-6.10 (m, 1H), 7.40 (d, 2H) and 7.50 (d, 2H).

MS: ESP+ (M+H) = 345.

5 **Example 50:** There is no compound with Example No. 50.

Example 51: N-((5S)-3-(4-(2-Benzoyloxymethyl-3,6- and -5,6-dihydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

Reference Example 21 (0.28 g), palladium(0) bis(dibenzylideneacetone) (0.039 g),
10 triphenylarsine (0.052 g) and lithium chloride (0.107 g) were dissolved in degassed *N,N*-dimethylformamide (8 ml). The stannane (Reference Example 1) (0.338 g) was added after 5 minutes and stirring was continued for 21 hours at 40 °C. Aqueous potassium fluoride (2.0 ml, 2M) was added and stirring continued for 20 minutes whereupon the mixture was evaporated. The residue was partitioned between ethyl acetate (40 ml) and water (10 ml), and the organic
15 layer dried and evaporated to an oil which was purified by silica-gel MPLC [using a mixture of 3% MeOH and CH₂Cl₂ as eluant]. The resultant foam was triturated with diethyl ether to give a mixture of the title compounds (0.204g, 58%).

NMR (300 MHz, CDCl₃) d : 2.01 (s, H), 2.29-2.73 (m, 2H), 3.54-3.94 (m, 6H), 4.06 (t, 1H), 4.14-4.23 (m, 0.5H), 4.39-4.53 (m, 1.5H), 4.56-4.70 (m, 2H), 4.71-4.84 (m, 1H), 6.07 (d, 1H),
20 6.15 (t, 1H) and 7.29-7.51 (m, 9H).

MS: ESP+ (M+H)⁺ = 437.

Example 51A: N-((5S)-3-(4-(2-Benzoyloxymethyl-3,6-dihydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

Example 51A was prepared in an identical fashion to Example 51, but starting from Reference
25 Example 24 instead of Reference Example 21.

NMR (300 MHz, CDCl₃) d: 2.01 (s, 3H), 2.28-2.50 (m, 2H), 3.54-3.94 (m, 6H), 4.09 (t, 1H), 4.32-4.50 (m, 2H), 4.61 (d, 1H), 4.67 (d, 1H), 4.72-4.83 (m, 1H), 6.10 (br s, 1H), 6.18 (t, 1H), 7.28-7.50 (m, 9H).

MS: ESP⁺ (M+H) = 437.

5

Example 52: N-((5S)-3-(4-(2-Hydroxymethyl-3,6- and -5,6-dihdropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

The benzyl ether Example 51 (0.158 g) in dry dichloromethane (7 ml) at 0°C was treated with boron trichloride-dimethyl sulfide (0.097 g, 0.27 ml) and stirred for 3 days. TLC indicated
10 consumption of starting material. Methanol (5 ml) was added and stirring continued for 20 minutes. The solution was evaporated and the residue purified by silica-gel MPLC [using a mixture of 2% MeOH and CH₂Cl₂ as eluant] to give a mixture of the title compounds (0.043g, 35%).

NMR (300 MHz, CDCl₃) d: 2.01 (s, 3H), 2.21-2.70 (m, 2 H), 3.55-3.85 (m, 6H), 4.09 (t, 15 1H), 4.10-4.20 (m, 0.5H), 4.35-4.45 (m, 1.5H), 4.70-4.82 (m, 1H), 5.98-6.15 (m, 2H) and 7.35-7.50 (m, 4H).

MS: (ESP⁺) (M+H) = 347.

Example 52A: N-((5S)-3-(4-(2-Hydroxymethyl-3,6-dihdropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

20 Example 52A was prepared in an identical fashion to Example 52, but starting from Example 51A rather than Example 51.

NMR (300 MHz, DMSO-D₆) d: 1.82 (s, 3H), 2.11-2.40 (m, 2H), 3.37-3.79 (m, 7H), 3.99-4.35 (m, 2H), 4.10 (t, 1H), 4.62-4.80 (m, 1H), 6.18-6.25 (m, 1H), 7.40-7.55 (m, 4H), 8.20 (t, 1H).

25 MS: ESP⁺ (M+H) = 347.

Example 53: N-((5S)-3-(4-(2-Acetamidomethyl-3,6- and -5,6-dihydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

Trimethyl phosphite (1 ml) was added slowly under argon to a stirred solution of the azide Reference example 23 (0.138 g) in dimethoxyethane (2 ml) at 50°C. After 2.25 hours, the solution was evaporated, redissolved in dimethoxyethane (2 ml) and treated with 6N hydrochloric acid (0.065 ml). The mixture was refluxed for 0.25 hours, cooled and stirred at ambient temperature for 0.5 hours. The solution was evaporated to the crude amine hydrochloride. The amine hydrochloride (0.22 mmol) in dimethoxyethane (2 ml) and water (0.5 ml) was taken to pH 9 with 2.5M sodium hydroxide solution and then treated with acetic anhydride (0.075 g). After 2.5 hours, the solution was evaporated and purified by Isolute silica-gel chromatography [using a mixture of 5% MeOH and CH₂Cl₂ as eluant] to give the title mixture of compounds (0.063g, 45%).

NMR (300 MHz, CDCl₃) d: 1.99, 2.02 and 2.03 (s/s/s, together 6H), 2.30-2.40 (m, 1H), 3.20-3.32 (m, 1H), 3.58-3.83 (m, 6H), 4.09 (t, 1H), 4.10-4.19 and 4.30-4.41 (m/m, together 2H), 4.74-4.82 (m, 1H), 5.98 and 6.09 (each broad s, together 2H), 6.43 (broad s, 1H), 7.32-7.40 and 7.43-7.50 (m/m, together 4H).

MS: ESP+ (M+H) = 388.

Example 54: N-((5S)-3-(4-(2-{Pyrimid-2-ylthio}methyl-3,6-dihydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

The alcohol Example 52A (0.162 g), 2-mercaptopyrimidine (0.105 g) and *N,N*-dimethylformamide dincopentyl acetal (0.326 g, 0.393 ml) in dry acetonitrile (10 ml) were refluxed for 4.5 hours. The solution was cooled, evaporated and purified by silica-gel MPLC [using a mixture of 6 % MeOH and CH₂Cl₂ as eluant] to give an oil which was triturated with diethyl ether to afford the title compound in crystalline form (0.134g, 63%).

NMR (300 MHz, CDCl₃) d: 2.02 (s, 3H), 2.30-2.72 (m, 2H), 3.39-4.00 (m, 6H), 4.05 (t, 1H), 4.13-4.62 (m, 2H), 4.72-4.82 (m, 1H), 6.09-6.25 (m, 2H), 6.98 (t, 1H), 7.40 (d, 2H), 7.49 (d, 2H), 8.52 (d, 2H).

MS: ESP+ (M+H) = 441.

Examples 55 to 98 : Carboxamides


Using a similar procedure to that described in Example 35, but starting with the appropriate
5 carboxylic acid instead of 4-tert-butylhexanecarboxylic acid the following compounds of
the formula XIV (Table A) were prepared by robotic synthesis (using a robot machine
manufactured by Zymark). The procedure used is described below.

The TFA salt of Example 2 in DMF (0.75mM, 4ml) was added to the appropriate carboxylic
10 acid (0.863mM) under argon. To this was then added a solution of N-methylmorpholine in
DMF (1mM, 1ml) and a solution of N-hydroxybenzotriazole in DMF (0.863mM, 1ml). The
mixture was then stirred at ambient temperature for 5 minutes after which time a solution of
1-ethyl-3-dimethylaminopropyl-carbodiimide (EDC) in DMF (0.863mmole, 3ml) was added.
The reaction was then stirred at ambient temperature overnight. The solvent was evaporated
15 using centrifugal evaporation (SAVANT AES2000) using radiant cover for 3 hours. The
residue was dissolved in CH₂Cl₂ (5ml). The organics were sequentially washed with 1N HCl
(4ml), water (4ml), saturated NaHCO₃ (4ml) and brine (5ml). The solvent was again removed
by centrifugal evaporation (SAVANT AES2000) using low heat for 3 hours and the products
transferred into 16x49mm vials.

20 The less pure products were subjected to further purification by Bondelut Chromatography
using conditions comparable to those of the TLC system used in earlier Examples ie. 5-10%
MeOH / CH₂Cl₂. The relevant fractions were combined and the solvent removed by
centrifugal evaporation (SAVANT AES2000) on low heat for 3 hours. The products were
then transferred into 16x49mm vials.


25

The compounds were analysed by mass spectroscopy and were characterised by HPLC
retention time, using the HPLC conditions described in Example 35.

Table A : N-Acyl compounds:

(XIV)

Example No.	R (Formula (XIV))	MS Mol Ion (M+H)	Hplc retention time (min.)
55	[1R-(1A,2B,3A)]-(+)-3-methyl-2-(nitromethyl)-5-oxocyclopentaneacetyl-	513	19.17
56	2-butoxyacetyl-	430	19.57
57	5-fluoroindole-3-acetyl-	491	19.78
58	3-acetyl-2,2-dimethylcyclobutyl-acetyl-	482	19.17
59	2-cyclopenten-1-acetyl-	424	20.15
60	2-naphthoxyacetyl-	500	21.63
61	oxamoyl-	463	19.95
62	N-t-butoxycarbonylglycyl-	473	19.08
63	methylthioacetyl-	404	17.77
64	There is no compound with Example No.64		
65	N-methyl-2-pyrroloyl-	423	19.12
66	trifluoroacetyl	412	20.12
67	3-(2-furyl)-acryloyl-	436	19.60
68	3-furoyl-	410	18.10
69	3-(2-thienyl)-acryoyl-	452	20.23
70	thiophene-2-acetyl-	440	19.28
71	thiophen-3-oyl-	426	18.78
72	There is no compound with Example No. 72		
73	indole-3-acetyl-	473	19.50
74	There is no compound with Example No. 74		
75	There is no compound with Example No. 75		
76	(2-pyrimidylthio)-acetyl-	468	17.68

Table A continued : N-Acyl compounds:

Example No.	R (Formula (XIV))	MS Mol Ion (M+H)	Hplc retention time (min.)
77	3,4-dihydro-2H-pyranoyl-	426	18.77
78	5-(O-methyl)-comenoyl	468	15.15
79	1,4-benzodioxan-6-acetyl-	492	19.32
80	3-(phenylsulfonyl)-propionoyl-	512	18.88
81	4-(aminosulfonyl)-benzoyl-	499	16.93
82	4-(diethylaminosulfonyl)-benzoyl-	583	22.38
83	4-(dimethylamino)-butyroyl-	429	15.20
84	N-acetyl-L-prolyl-	455	16.23
85	2,3,4-trimethoxycinnamoyl-	536	20.42
86	There is no compound with Example No. 86		
87	(R)-5-oxo-2-tetrahydrofuroyl-	428	16.43
88	7-oxo-octanoyl-	456	18.38
89	There is no compound with Example No. 89		
90	2-benzofuranoyl-	460	20.95
91	indole-1-acetyl-	473	20.97
92	sulfolanyl-3-acetyl-	476	16.67
93	4-oxo-4H-1-benzo-2-pyranoyl-	488	19.18
94	There is no compound with Example No. 94		
95	indole-3-oyl-	459	18.98
96	2-(n-propylthio)-nicotinoyl-	495	20.62
97	2-methoxy-pyridin-3-oyl-	451	17.95
98	2,3-dihydro-5-oxothiazolo-[3,2-A]-pyrimidin-6-oyl-	496	16.37


Examples 99 TO 139: Sulfonamides

Using a similar procedure to that described in Example 38, but starting with the appropriate sulfonyl chloride instead of p-toluenesulfonyl chloride the following compounds of the formula XV (Table B) were prepared by robotic synthesis (using a robot machine
5 manufactured by Zymark) . The procedure used is described below.

The TFA salt of Example 2 (279mg, 0.65mM) was suspended in CH_2Cl_2 (10ml). To this was added triethylamine (243 μl) and then a solution of the appropriate sulfonyl chloride in CH_2Cl_2 (0.78mM in 5ml). The mixture went into solution after 10-15 minutes and the reaction was
10 then stirred at ambient temperature for a further 4 hours. The compounds were subjected to purification by Bondelut Chromatography using conditions comparable to those of the TLC system used in earlier Examples ie. 5-10% MeOH / CH_2Cl_2 . The relevant fractions were combined and the solvent removed by centrifugal evaporation (SAVANT AES2000) on low heat for 3 hours. The products were then transferred into 16x49mm vials.


15

The compounds were analysed by mass spectroscopy and were characterised by HPLC retention time, using the HPLC conditions described in Example 35.

Table B : N-Sulfonyl compounds:

Example No.	R (Formula (XV))	MS Mol Ion (M+H)	Hplc retention time (min.)
99	8-quinoline-sulfonyl-	507	19.93
100	2-naphthalene-sulfonyl-	504	22.82
101	2-methoxycarbonylphenyl-sulfonyl	514	20.65
102	4-fluorobenzene-sulfonyl-	474	21.38
103	3-chloropropane-sulfonyl-	456	20.17
114	4-chlorobenzene-sulfonyl-	490	22.43
115	4-carboxybenzene-sulfonyl-	498*	19.75
116	isopropylsulfonyl-	422	19.28
117	b-styrylsulfonyl-	482	21.88
118	ethanesulfonyl-	408	18.35
119	1-butanesulfonyl-	436	20.70
120	1-octanesulfonyl-	492	25.07
121	2-chlorobenzene-sulfonyl-	490	21.67
122	3-carboxybenzene-sulfonyl-	498*	19.23
123	3-trifluoromethylbenzene-sulfonyl-	524	22.77
124	2,2,2-trifluorethanesulfonyl-	462	20.48
125	5-chlorothiophene-sulfonyl-	496	22.72
126	2-(pyrid-2-yl)-thiophene-5-sulfonyl-	539	21.68
127	2,4-difluorobenzene-sulfonyl-	492	21.62
128	3-fluorobenzene-sulfonyl-	474	21.50
129	4-acetamidobenzene-sulfonyl-	513	18.90
130	1-methylimidazol-4-yl-sulfonyl-	460	17.03

* M-H

Table B continued : N-Sulfonyl compounds:

Example No.	R (Formula (XV))	MS Mol Ion (M+H)	Hplc retention time (min.)
131	5-(isoxazol-3-yl)-thiophene-2-sulfonyl-	529	21.50
132	5-bromothiophene-2-sulfonyl-	540	22.83
134	4-cyanobenzene-sulfonyl-	481	20.85
135	4-trifluoromethoxybenzene-sulfonyl-	540	22.15
136	4-trifluoromethylbenzene-sulfonyl-	524	22.93
137	3,4-dimethoxybenzene-sulfonyl-	516	20.27
138	5-dimethylamino-1-naphthalene-sulfonyl-	549	20.15
139	benzenesulfonyl-	456	21.00

NB. There are no compounds with Example Nos. 104-113 and 133.

5 **Example 140**

The following illustrate representative pharmaceutical dosage forms containing the compound of formula I, or a pharmaceutically-acceptable salt thereof (hereafter compound X), for therapeutic or prophylactic use in humans:

10 (a)	Tablet I	mg/tablet
	Compound X.....	100
	Lactose Ph.Eur.....	179
	Croscarmellose sodium.....	12
	Polyvinylpyrrolidone.....	6
15	Magnesium stearate.....	3

(b)	<u>Tablet II</u>	<u>mg/tablet</u>
	Compound X.....	50
	Lactose Ph.Eur.....	229
	Croscarmellose sodium.....	12
5	Polyvinylpyrrolidone.....	6
	Magnesium stearate.....	3
(c)	<u>Tablet III</u>	<u>mg/tablet</u>
	Compound X.....	1
10	Lactose Ph.Eur.....	92
	Croscarmellose sodium.....	4
	Polyvinylpyrrolidone.....	2
	Magnesium stearate.....	1
15 (d)	<u>Capsule</u>	<u>mg/capsule</u>
	Compound X.....	10
	Lactose Ph.Eur	389
	Croscarmellose sodium.....	100
	Magnesium stearate	1
20	(e)	<u>Injection I</u>
		<u>(50 mg/ml)</u>
	Compound X	5.0% w/v
	Isotonic aqueous solution	to 100%


25 Buffers, pharmaceutically-acceptable cosolvents such as polyethylene glycol, polypropylene glycol, glycerol or ethanol or complexing agents such as hydroxy-propyl β cyclodextrin may be used to aid formulation.

Note

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate.

CLAIMS

1. A compound of the formula (I):



(I)


wherein:

- 10 R^1 is hydroxy, chloro, fluoro, (1-4C)alkanesulfonyloxy, amino, azido, (1-4C)alkoxy, (1-4C)alkylthio, (1-4C)alkylaminocarbonyloxy;
or of the formula $-NHC(=O)R^b$ wherein R^b is hydrogen, (1-4C)alkoxy, amino, chloromethyl, dichloromethyl, cyanomethyl, methoxymethyl, acetylmethyl, methylamino, dimethylamino or (1-4C)alkyl;
- 15 or R^1 is of the formula $-NHS(O)_n(1-4C)alkyl$ wherein n is 0, 1 or 2;
 R^2 and R^3 are independently hydrogen or fluoro;
 D is O, S, SO, SO_2 or NR^7 ;
when D is O, R^4 and R^5 are independently hydroxy, bromo, oxo ($=O$), (1-4C)alkyl, (1-4C)alkanoylamino-(1-4C)alkyl, hydroxy-(1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl,
- 20 AR-oxymethyl, AR-thiomethyl (wherein AR is as defined hereinbelow) or independently as defined for R^7 hereinbelow;
when D is S, SO, SO_2 or NR^7 , R^4 and R^5 are independently oxo ($=O$), (1-4C)alkyl, (1-4C)alkanoylamino-(1-4C)alkyl, hydroxy-(1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl, AR-oxymethyl, AR-thiomethyl (wherein AR is as defined
- 25 hereinbelow) or independently as defined for R^7 hereinbelow;
 R^6 is hydrogen, (1-4C)alkyl, hydroxy, (1-4C)alkoxy or (2-4C)alkanoyloxy;
 $>A-B-$ is of the formula $>C=C(R^a)-$, $>CHCHR^a-$, or $>C(OH)CHR^a-$ ($>$ represents two single bonds) wherein R^a is hydrogen or (1-4C)alkyl;

R^7 is hydrogen, cyano, 2-((1-4C)alkoxycarbonyl)ethenyl, 2-cyanoethenyl,
 2-cyano-2-((1-4C)alkyl)ethenyl, 2-((1-4C)alkylaminocarbonyl)ethenyl, AR (as defined
 hereinbelow) or a tetrazole ring system (optionally mono-substituted in the 1- or 2- position of
 the tetrazole ring) wherein the tetrazole ring system is joined to the nitrogen in NR^7 by a ring
 5 carbon atom;
 or R^7 is of the formula $R^{10}CO-$, $R^{10}SO_2-$ or $R^{10}CS-$
 wherein R^{10} is AR (as defined hereinbelow), cyclopentyl or cyclohexyl (wherein the last two-
 mentioned cycloalkyl rings are optionally mono- or disubstituted by substituents
 independently selected from (1-4C)alkyl (including geminal disubstitution), hydroxy,
 10 (1-4C)alkoxy, (1-4C)alkylthio, acetamido, (1-4C)alkanoyl, cyano and trifluoromethyl),
 (1-4C)alkoxycarbonyl, hydrogen, amino, trifluoromethyl, (1-4C)alkylamino,
 di((1-4C)alkyl)amino, 2,3-dihydro-5-oxothiazolo-[3,2-A]pyrimidin-6-yl, 2-(2-furyl)ethenyl,
 2-(2-thienyl)ethenyl, 2-phenylethenyl (wherein the phenyl substituent is optionally substituted
 by up to three substituents independently selected from (1-4C)alkoxy, halo and cyano), 3,4-
 15 dihydropyran-2-yl, coumal-5-yl, 5-methoxy-4-oxopyran-2-yl, N-acetylpyrrolidin-2-yl, 5-oxo-
 tetrahydrofuran-2-yl, benzopyranone or (1-10C)alkyl [wherein (1-10C)alkyl is optionally
 substituted by hydroxy, cyano, halo, (1-10C)alkoxy, trifluoromethyl,
 (1-4C)alkoxy-(1-4C)alkoxy, (1-4C)alkoxy-(1-4C)alkoxy-(1-4C)alkoxy, (1-4C)alkanoyl,
 (1-4C)alkoxycarbonyl, amino, (1-4C)alkylamino, di((1-4C)alkyl)amino, (1-
 20 6C)alkanoylamino, (1-4C)alkoxycarbonylamino, \underline{N} -(1-4C)alkyl- \underline{N} -(2-6C)alkanoylamino, (1-
 4C)alkylS(O)_pNH-,
 (1-4C)alkylS(O)_p((1-4C)alkyl)N-, fluoro(1-4C)alkylS(O)_pNH-,
 fluoro(1-4C)alkylS(O)_p((1-4C)alkyl)N-, phosphono, (1-4C)alkoxy(hydroxy)phosphoryl,
 di-(1-4C)alkoxyphosphoryl, (1-4C)alkylS(O)_q-, phenylS(O)_q- (wherein the phenyl group is
 25 optionally substituted by up to three substituents independently selected from (1-4C)alkoxy,
 halo and cyano), or CY (as defined hereinbelow), wherein p is 1 or 2 and q is 0, 1 or 2];
 or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is an optionally substituted 5- or
 6-membered heteroaryl, optionally substituted phenyl, (1-4C)alkylamino,
 benzyloxy-(1-4C)alkyl or optionally substituted (1-10C)alkyl;
 30 or R^{10} is of the formula $R^{12}O-$ wherein R^{12} is optionally substituted (1-6C)alkyl;

- or R⁷ is of the formula R^dOC(R^e)=CH(C=O)-, R^fC(=O)C(=O)-, R^gN=C(R^h)C(=O)- or RⁱNHC(R^j)=CHC(=O)- wherein R^d is (1-6C)alkyl, R^e is hydrogen or (1-6C)alkyl, or R^d and R^e together form a (3-4C)alkylene chain, R^f is hydrogen, (1-6C)alkyl, hydroxy(1-6C)alkyl, (1-6C)alkoxy(1-6C)alkyl, amino, (1-4C)alkylamino, 5 di-(1-4C)alkylamino, (1-6C)alkoxy, (1-6C)alkoxy(1-6C)alkoxy, hydroxy(2-6C)alkoxy, (1-4C)alkylamino(2-6C)alkoxy, di-(1-4C)alkylamino(2-6C)alkoxy, R^g is (1-6C)alkyl, hydroxy or (1-6C)alkoxy, R^h is hydrogen or (1-6C)alkyl, Rⁱ is hydrogen, (1-6C)alkyl, optionally substituted phenyl or an optionally substituted 5- or 6-membered heteroaryl and R^j is hydrogen or (1-6C)alkyl;
- 10 or R⁷ is of the formula R¹⁴CH(R¹³)(CH₂)_m - wherein m is 0 or 1, R¹³ is fluoro, cyano, (1-4C)alkoxy, (1-4C)alkylsulfonyl, (1-4C)alkoxycarbonyl or hydroxy, (provided that when m is 0, R¹³ is not fluoro or hydroxy) and R¹⁴ is hydrogen or (1-4C)alkyl; wherein AR is optionally substituted phenyl, optionally substituted phenyl(1-4C)alkyl, optionally substituted 5- or 6-membered heteroaryl, optionally substituted naphthyl or an 15 optionally substituted 5/6 or 6/6 bicyclic heteroaryl ring system, in which the bicyclic heteroaryl ring systems may be linked via an atom in either of the rings comprising the bicyclic system, and wherein the mono- and bicyclic heteroaryl ring systems are linked via a ring carbon atom; wherein CY is a 4-, 5- or 6-membered cycloalkyl ring, a 5- or 6-membered cycloalkenyl ring, 20 naphthoxy, thiophen-2-yl, indol-1-yl, indol-3-yl, pyrimidin-2-ylthio, 1,4-benzodioxan-6-yl, sulfolan-3-yl, pyridin-2-yl; wherein any of the afore-mentioned ring systems in CY may be optionally substituted by up to three substituents independently selected from halo, (1-4C)alkyl (including geminal disubstitution when CY is a cycloalkyl or cycloalkenyl ring), acyl, oxo and nitro-(1-4C)alkyl; and pharmaceutically-acceptable salts 25 thereof; except that N-((5S)-3-(4-(2-oxo-5,6-dihydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide is excluded.
2. A compound of the formula (I) as claimed in claim 1, except that in the definition of R⁴ and R⁵, neither R⁴ nor R⁵ can be oxo (=O).
- 30 3. A compound, as claimed in claims 1 and 2, of the formula (IB):

- 88 -




(IB)

wherein R¹ is hydroxy or acetamido; R² is hydrogen or fluoro and R⁴ and R⁵ are independently
 5 hydrogen, hydroxy, bromo, (1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl, hydroxymethyl, (1-4C)alkoxymethyl or carbamoyl; and pharmaceutically-acceptable salts thereof.

4. A compound, as claimed in claims 1 and 2, of the formula (IB) wherein R¹ is acetamido; R² is hydrogen or fluoro and R⁴ and R⁵ are independently hydrogen, AR-oxymethyl or AR-thiomethyl (wherein AR is phenyl, phenyl(1-4C)alkyl, naphthyl, furan,
 10 pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole or thiophene); and pharmaceutically-acceptable salts thereof.

5. A compound, as claimed in claims 1 and 2, of the formula (IC):



(IC)

15

wherein R¹ is hydroxy or acetamido; R² is hydrogen or fluoro; R⁴ and R⁵ are independently
 hydrogen, AR-oxymethyl or AR-thiomethyl (wherein AR is phenyl, phenyl(1-4C)alkyl,
 naphthyl, furan, pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine,
 20 isoxazole, oxazole, isothiazole, thiazole or thiophene), hydroxy, bromo, (1-4C)alkyl, carboxy,
 (1-4C)alkoxycarbonyl, hydroxymethyl, (1-4C)alkoxymethyl or carbamoyl and R⁷ is cyano,
 pyrimidin-2-yl, tetrazol-5-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl or R⁷ is of the

- formula $R^{10}CO-$ or $R^{10}SO_2-$ (wherein R^{10} is hydrogen, (1-4C)alkyl [optionally substituted by hydroxy, (1-4C)alkoxy, (1-4C)alkylS(O)_q, (1-4C)alkylamino, (1-4C)alkanoyl, naphthoxy, (2-6C)alkanoylamino or (1-4C)alkylS(O)_pNH- wherein p is 1 or 2 and q is 0, 1 or 2], phenyl, naphthyl, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole, thiophene, benzofuran, benzoimidazole, benzothiophene, benzisothiazole, benzoxazole, benzisoxazole, pyridoimidazole, pyrimidoimidazole, quinoline, quinoxaline, quinazoline, phthalazine, cinnoline or naphthyridine, or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is (1-6C)alkyl), or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is (1-6C)alkoxy; and pharmaceutically-acceptable salts thereof.
- 10 6. A compound, as claimed in claims 1, 2 and 5, of the formula (IC), wherein R^1 is acetamido; R^2 is hydrogen or fluoro; R^4 and R^5 are hydrogen; and R^7 is pyrimidin-2-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl, or R^7 is of the formula $R^{10}CO-$ (wherein R^{10} is hydrogen or (1-4C)alkyl [optionally substituted by hydroxy or (1-4C)alkylS(O)_q wherein q is 0, 1 or 2], or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is (1-6C)alkyl), or R^7 is
 15 of the formula $R^fC(=O)C(=O)-$ wherein R^f is (1-6C)alkoxy; and pharmaceutically-acceptable salts thereof.
7. A compound, as claimed in claims 1, 2 and 5, of the formula (IC), wherein R^1 is acetamido; R^2 is hydrogen or fluoro; R^4 and R^5 are hydrogen; and R^7 is pyrimidin-2-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl, or R^7 is of the formula $R^{10}SO_2-$ (wherein R^{10} is
 20 hydrogen or (1-4C)alkyl [optionally substituted by hydroxy or (1-4C)alkylS(O)_q wherein q is 0, 1 or 2], or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is (1-6C)alkyl), or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is (1-6C)alkoxy; and pharmaceutically-acceptable salts thereof.
8. A compound of the formula (I), or a pharmaceutically-acceptable salt thereof, as
 25 claimed in claims 1 and 2, selected from :-
- N-((5S)-3-(4-(1-methylsulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxo-oxazolidin-5-ylmethyl)acetamide;
- N-((5S)-3-(4-(1-acetyloxymethylcarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
- 30 N-((5S)-3-(4-(1-hydroxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

- 90 -

N-((5S)-3-(3-fluoro-4-(-acetyloxymethylcarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(3-fluoro-4-(1-hydroxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

5 N-((5S)-3-(4-(2,3-dihydro-6H-pyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(3-fluoro-4-(2,3-dihydro-6H-pyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(3-fluoro-4-(1-{pyrimid-2-yl}-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

10 N-((5S)-3-(3-fluoro-4-(1-methylsulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(4-(1-{pyrimid-2-yl}-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(3-fluoro-4-(1-acetoacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

15 N-((5S)-3-(4-(1-(naphth-2-ylsulfonyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;


N-((5S)-3-(4-(1-(naphth-2-oxyacetyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide; and

20 N-((5S)-3-(4-(1-(methylthioacetyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

9. A process for the preparation of a compound of the formula (I) as claimed in claim 1, which comprises :-

(a) the deprotection of a compound of formula (II):


25



(II)


- (b) the modification of a substituent in, or the introduction of a substituent into, another compound of formula (I) or (II);
- (c) when R^1 or R^{20} is of the formula $-NHS(O)_n(1-4C)alkyl$, wherein n is 1 or 2, the oxidation of a compound of the formula (I) or (II) wherein n is 0 or, when n is 2 the oxidation of a compound of the formula (I) or (II) wherein n is 1;
- (d) when R^1 or R^{20} is of the formula $-NHC(=O)R^b$ or $NHS(O)_n(1-4C)alkyl$, the introduction of the group $-C(=O)R^b$ or $-S(O)_n(1-4C)alkyl$ into a compound of the formula (III);

10




(III)

- (e) when R¹ or R²⁰ is hydroxy, the reaction of a compound of the formula (V) with a compound of formula (VI):



(M)




(VI)


20

- (f) when $>A1-B1-$ is $>C=CR^{a1-}$, the reaction of a compound of the formula (VII) with a compound of the formula (VIII):

- 92 -




(VII)



(VIII)


(g) when >A1-B1- is >CHCH(R^{a1})-, the catalytic hydrogenation of a compound of the formula (I) or (II) wherein >A1-B1- is >C=CR^{a1}-;

(h) when >A1-B1- is >C=CR^{a1}-, the elimination of the elements of water, or HOCOR²³ or HOSO₂ R²⁴ from a compound of the formula (IX) (ie. when R²⁵ is -H, -COR²³ or -SO₂ R²⁴);



(IX)

(i) when D is NR⁷ and R⁷ is R¹⁰CO- or R¹⁰S(O)_n-, wherein n is 2, the reaction of a compound of formula (X) with a compound of the formula (XI) or (XII), wherein n is 2:



(X)




(XI)



(XII)

15

- (j) when R^1 or R^{20} is azido, the reaction of a compound of the formula (XIII) with a source of azide:



5

(XIII)

- (k) when R^1 or R^{20} is amino, the reduction of a compound of the formula (I) or (II) wherein R^1 or R^{20} is azido;
- 10 (l) the preparation when R^1 or R^{20} is chloro, fluoro, (1-4C)alkanesulfonyloxy, or (1-4C)alkylaminocarbonyloxy, or R^{20} is of the formula $-N(CO_2R^{27})CO(1-4C)alkyl$ from a compound of the formula (I) and (II) wherein R^1 or R^{20} is hydroxy; or
- (m) the preparation when R^1 or R^{20} is chloro, (1-4C)alkylthio or (1-4C)alkoxy, from a compound of the formula (XIII);
- 15 wherein R^2 , R^3 and R^6 are as hereinabove defined; R^{18} is R^4 or protected R^4 ; R^{19} is R^5 or protected R^5 ; R^{20} is R^1 or protected R^1 ; $>A1-B1-$ is $>A-B-$ or protected $>A-B-$ and D1 is D in which functional groups are optionally protected; R^{21} is R^8 or protected R^8 ; R^{21} is (1-6C)alkyl or benzyl; R^{22} is of the formula (1-4C)alkyl or $-S(O)_n(1-4C)alkyl$; R^{23} is (1-4C)alkyl; R^{24} is an optionally substituted phenyl group; R^{25} is hydrogen, $-COR^{23}$
- 20 or $-SO_2R^{24}$; R^{26} is mesyloxy or tosyloxy; R^{27} is (1-4C)alkyl or benzyl; n is 0, 1 or 2 unless otherwise stated above; L^1 is an iodo or triflate leaving group; L^2 is a leaving group; and Y is a trialkyltin residue or a boronate acid or ester residue; and thereafter if necessary:
- removing any protecting groups;
 - forming a pharmaceutically-acceptable salt;
- 25 and when an optically active form of a compound of the formula (I) is required, it may be obtained by carrying out one of the above procedures using an optically active starting material, or by resolution of a racemic form of the compound or intermediate using a standard


procedure; and similarly, when a pure regioisomer of a compound of the formula (I) is required, it may be obtained by carrying out one of the above procedures using a pure regioisomer as a starting material, or by resolution of a mixture of the regioisomers or intermediates using a standard procedure.

- 5 10. A pharmaceutical composition which comprises a compound of the formula (I) or a pharmaceutically-acceptable salt thereof, as claimed in claims 1 to 8 and a pharmaceutically-acceptable diluent or carrier.
11. The use of a compound of the formula (I) or a pharmaceutically-acceptable salt thereof, as claimed in claims 1 to 8, in the manufacture of a novel medicament for use in
- 10 the production of an antibacterial effect in a warm blooded animal such as man.
12. A method for producing an antibacterial effect in a warm blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a compound of the present invention, or a pharmaceutically-acceptable salt thereof.

AMENDED CLAIMS

[received by the International Bureau on 24 July 1997 (24.07.97);
original claims 1,3 and 5-12 amended; remaining claims unchanged (11 pages)]

1. A compound of the formula (I):



5

(I)

wherein:

- R¹ is hydroxy, chloro, fluoro, (1-4C)alkanesulfonyloxy, amino, azido, (1-4C)alkoxy,
10 (1-4C)alkylthio, (1-4C)alkylaminocarbonyloxy;

or of the formula -NHC(=O)R^b wherein R^b is hydrogen, (1-4C)alkoxy, amino, chloromethyl, dichloromethyl, cyanomethyl, methoxymethyl, acetylmethyl, methylamino, dimethylamino or (1-4C)alkyl;

or R¹ is of the formula -NHS(O)_n(1-4C)alkyl wherein n is 0, 1 or 2;

- 15 R² and R³ are independently hydrogen or fluoro;

D is O, S, SO, SO₂ or NR⁷;

when D is O, R⁴ and R⁵ are independently hydroxy, bromo, oxo (=O), (1-4C)alkyl, (1-4C)alkanoylamino-(1-4C)alkyl, hydroxy-(1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl, AR-oxymethyl, AR-thiomethyl (wherein AR is as defined

- 20 hereinbelow) or independently as defined for R⁷ hereinbelow;

when D is S, SO, SO₂ or NR⁷, R⁴ and R⁵ are independently oxo (=O), (1-4C)alkyl, (1-4C)alkanoylamino-(1-4C)alkyl, hydroxy-(1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl, AR-oxymethyl, AR-thiomethyl (wherein AR is as defined hereinbelow) or independently as defined for R⁷ hereinbelow;

- 25 R⁶ is hydrogen, (1-4C)alkyl, hydroxy, (1-4C)alkoxy or (2-4C)alkanoyloxy;

- >A-B- is of the formula $>C=C(R^a)-$, $>CHCHR^a-$ or $>C(OH)CHR^a-$ (> represents two single bonds) wherein R^a is hydrogen or (1-4C)alkyl;
- R^7 is hydrogen, cyano, 2-((1-4C)alkoxycarbonyl)ethenyl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl, 2-((1-4C)alkylaminocarbonyl)ethenyl, AR (as defined hereinbelow) or a tetrazole ring system (optionally mono-substituted in the 1- or 2- position of the tetrazole ring) wherein the tetrazole ring system is joined to the nitrogen in NR^7 by a ring carbon atom;
- or R^7 is of the formula $R^{10}CO-$, $R^{10}SO_2-$ or $R^{10}CS-$ wherein R^{10} is AR (as defined hereinbelow), cyclopentyl or cyclohexyl (wherein the last two-mentioned cycloalkyl rings are optionally mono- or disubstituted by substituents independently selected from (1-4C)alkyl (including geminal disubstitution), hydroxy, (1-4C)alkoxy, (1-4C)alkylthio, acetamido, (1-4C)alkanoyl, cyano and trifluoromethyl),
- (1-4C)alkoxycarbonyl, hydrogen, amino, trifluoromethyl, (1-4C)alkylamino, di((1-4C)alkyl)amino, 2,3-dihydro-5-oxothiazolo-[3,2-A]pyrimidin-6-yl, 2-(2-furyl)ethenyl, 2-(2-thienyl)ethenyl, 2-phenylethenyl (wherein the phenyl substituent is optionally substituted by up to three substituents independently selected from (1-4C)alkoxy, halo and cyano), 3,4-dihydropyran-2-yl, coumal-5-yl, 5-methoxy-4-oxopyran-2-yl, N-acetylpyrrolidin-2-yl, 5-oxo-tetrahydrofuran-2-yl, benzopyranone or (1-10C)alkyl [wherein (1-10C)alkyl is optionally substituted by hydroxy, cyano, halo, (1-10C)alkoxy, trifluoromethyl, (1-4C)alkoxy-(1-4C)alkoxy, (1-4C)alkoxy-(1-4C)alkoxy-(1-4C)alkoxy, (1-4C)alkanoyl, (1-4C)alkoxycarbonyl, amino, (1-4C)alkylamino, di((1-4C)alkyl)amino, (1-6C)alkanoylamino,
- (1-4C)alkoxycarbonylamino, \underline{N} -(1-4C)alkyl- \underline{N} -(2-6C)alkanoylamino, (1-4C)alkylS(O)_pNH-, (1-4C)alkylS(O)_p((1-4C)alkyl)N-, fluoro(1-4C)alkylS(O)_pNH-, fluoro(1-4C)alkylS(O)_p((1-4C)alkyl)N-, phosphono, (1-4C)alkoxy(hydroxy)phosphoryl, di-(1-4C)alkoxyphosphoryl, (1-4C)alkylS(O)_q-, phenylS(O)_q- (wherein the phenyl group is optionally substituted by up to three


- substituents independently selected from (1-4C)alkoxy, halo and cyano), or CY (as defined hereinbelow), wherein p is 1 or 2 and q is 0, 1 or 2] ;
- or R¹⁰ is of the formula R¹¹C(O)O(1-6C)alkyl wherein R¹¹ is an optionally substituted 5- or 6-membered heteroaryl, optionally substituted phenyl, (1-4C)alkylamino,
- 5 benzyloxy-(1-4C)alkyl or optionally substituted (1-10C)alkyl;
- or R¹⁰ is of the formula R¹²O- wherein R¹² is optionally substituted (1-6C)alkyl;
- or R⁷ is of the formula R^dOC(R^e)=CH(C=O)-, R^fC(=O)C(=O)-, R^gN=C(R^h)C(=O)- or RⁱNHC(R^j)=CHC(=O)- wherein R^d is (1-6C)alkyl, R^e is hydrogen or (1-6C)alkyl,
- or R^d and R^e together form a (3-4C)alkylene chain, R^f is hydrogen, (1-6C)alkyl,
- 10 hydroxy(1-6C)alkyl, (1-6C)alkoxy(1-6C)alkyl, amino, (1-4C)alkylamino, di-(1-4C)alkylamino, (1-6C)alkoxy, (1-6C)alkoxy(1-6C)alkoxy, hydroxy(2-6C)alkoxy, (1-4C)alkylamino(2-6C)alkoxy, di-(1-4C)alkylamino(2-6C)alkoxy, R^g is (1-6C)alkyl, hydroxy or (1-6C)alkoxy, R^h is hydrogen or (1-6C)alkyl,
- 15 Rⁱ is hydrogen, (1-6C)alkyl, optionally substituted phenyl or an optionally substituted 5- or 6-membered heteroaryl and R^j is hydrogen or (1-6C)alkyl;
- or R⁷ is of the formula R¹⁴CH(R¹³)(CH₂)_m - wherein m is 0 or 1, R¹³ is fluoro, cyano, (1-4C)alkoxy, (1-4C)alkylsulfonyl, (1-4C)alkoxycarbonyl or hydroxy, (provided that when m is 0, R¹³ is not fluoro or hydroxy) and R¹⁴ is hydrogen or (1-4C)alkyl;
- 20 wherein AR is optionally substituted phenyl, optionally substituted phenyl(1-4C)alkyl, optionally substituted 5- or 6-membered heteroaryl, optionally substituted naphthyl or an optionally substituted 5/6 or 6/6 bicyclic heteroaryl ring system, in which the bicyclic heteroaryl ring systems may be linked via an atom in either of the rings comprising the bicyclic system, and wherein the mono- and bicyclic heteroaryl
- 25 ring systems are linked via a ring carbon atom;
- wherein CY is a 4-, 5- or 6-membered cycloalkyl ring, a 5- or 6-membered cycloalkenyl ring, naphthoxy, thiophen-2-yl, indol-1-yl, indol-3-yl, pyrimidin-2-ylthio.
- 1,4-benzodioxan-6-yl, sulfolan-3-yl, pyridin-2-yl; wherein any of the afore-mentioned
- 30 ring systems in CY may be optionally substituted by up to three substituents

independently selected from halo, (1-4C)alkyl (including geminal disubstitution when CY is a cycloalkyl or cycloalkenyl ring), acyl, oxo and nitro-(1-4C)alkyl; pharmaceutically-acceptable salts thereof; and in-vivo-hydrolysable esters thereof; except that

5 N-((5S)-3-(4-(2-oxo-5,6-dihydropyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide is excluded.

2. A compound of the formula (I) as claimed in claim 1, except that in the definition of R^4 and R^5 , neither R^4 nor R^5 can be oxo (=O).

3. A compound, as claimed in claims 1 and 2, of the formula (IB):



10

(IB)


wherein R^1 is hydroxy or acetamido; R^2 is hydrogen or fluoro and R^4 and R^5 are independently hydrogen, hydroxy, bromo, (1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl, hydroxymethyl, (1-4C)alkoxymethyl or carbamoyl;

15 pharmaceutically-acceptable salts thereof; and in-vivo-hydrolysable esters thereof.

4. A compound, as claimed in claims 1 and 2, of the formula (IB) wherein R^1 is acetamido; R^2 is hydrogen or fluoro and R^4 and R^5 are independently hydrogen, AR-oxymethyl or AR-thiomethyl (wherein AR is phenyl, phenyl(1-4C)alkyl, naphthyl, furan, pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole or thiophene); and pharmaceutically-acceptable salts thereof.

20

5. A compound, as claimed in claims 1 and 2, of the formula (IC):



(IC)

- wherein R^1 is hydroxy or acetamido; R^2 is hydrogen or fluoro; R^4 and R^5 are independently hydrogen, AR-oxyethyl or AR-thiomethyl (wherein AR is phenyl, 5 phenyl(1-4C)alkyl, naphthyl, furan, pyrrole, pyrazole, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole or thiophene), hydroxy, bromo, (1-4C)alkyl, carboxy, (1-4C)alkoxycarbonyl, hydroxyethyl, (1-4C)alkoxyethyl or carbamoyl and R^7 is cyano, pyrimidin-2-yl, tetrazol-5-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl or R^7 is of
- 10 the formula $R^{10}CO-$ or $R^{10}SO_2-$ (wherein R^{10} is hydrogen, (1-4C)alkyl [optionally substituted by hydroxy, (1-4C)alkoxy, (1-4C)alkylS(O)_q, (1-4C)alkylamino, (1-4C)alkanoyl, naphthoxy, (2-6C)alkanoylamino or (1-4C)alkylS(O)_pNH- wherein p is 1 or 2 and q is 0, 1 or 2], phenyl, naphthyl, imidazole, triazole, pyrimidine, pyridazine, pyridine, isoxazole, oxazole, isothiazole, thiazole, thiophene, benzofuran, 15 benzoimidazole, benzothiophene, benzisothiazole, benzoxazole, benzisoxazole, pyridoimidazole, pyrimidoimidazole, quinoline, quinoxaline, quinazoline, phthalazine, cinnoline or naphthyridine, or R^{10} is of the formula $R^{11}C(O)O(1-6C)alkyl$ wherein R^{11} is (1-6C)alkyl), or R^7 is of the formula $R^fC(=O)C(=O)-$ wherein R^f is (1-6C)alkoxy; pharmaceutically-acceptable salts thereof; and in-vivo-hydrolysable esters
- 20 thereof.

6. A compound, as claimed in claims 1, 2 and 5, of the formula (IC), wherein R^1 is acetamido; R^2 is hydrogen or fluoro; R^4 and R^5 are hydrogen; and R^7 is pyrimidin-2-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl, or R^7 is of the formula $R^{10}CO-$ (wherein R^{10} is hydrogen or (1-4C)alkyl [optionally substituted by hydroxy or

(1-4C)alkylS(O)_q wherein q is 0, 1 or 2], or R¹⁰ is of the formula R¹¹C(O)O(1-6C)alkyl wherein R¹¹ is (1-6C)alkyl, or R⁷ is of the formula R^fC(=O)C(=O)- wherein R^f is (1-6C)alkoxy; pharmaceutically-acceptable salts thereof; and in-vivo-hydrolysable esters thereof.

7. A compound, as claimed in claims 1, 2 and 5, of the formula (IC), wherein R¹ is acetamido; R² is hydrogen or fluoro; R⁴ and R⁵ are hydrogen; and R⁷ is pyrimidin-2-yl, 2-cyanoethenyl, 2-cyano-2-((1-4C)alkyl)ethenyl, or R⁷ is of the formula R¹⁰SO₂- (wherein R¹⁰ is hydrogen or (1-4C)alkyl [optionally substituted by hydroxy or (1-4C)alkylS(O)_q wherein q is 0, 1 or 2], or R¹⁰ is of the formula R¹¹C(O)O(1-6C)alkyl wherein R¹¹ is (1-6C)alkyl, or R⁷ is of the formula R^fC(=O)C(=O)- wherein R^f is (1-6C)alkoxy; pharmaceutically-acceptable salts thereof; and in-vivo-hydrolysable esters thereof.

8. A compound of the formula (I), or a pharmaceutically-acceptable salt thereof, or an in-vivo-hydrolysable ester thereof when there is a suitable hydroxy group in the compound, as claimed in claims 1 and 2, selected from :-

N-((5S)-3-(4-(1-methylsulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxo-oxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(4-(1-acetyloxymethylcarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(4-(1-hydroxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(3-fluoro-4-(1-acetyloxymethylcarbonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

N-((5S)-3-(3-fluoro-4-(1-hydroxyacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;


N-((5S)-3-(4-(2,3-dihydro-6H-pyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;

- N-((5S)-3-(3-fluoro-4-(2,3-dihydro-6H-pyran-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
 N-((5S)-3-(3-fluoro-4-(1-{pyrimid-2-yl}-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
 5 N-((5S)-3-(3-fluoro-4-(1-methylsulfonyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
 N-((5S)-3-(4-(1-{pyrimid-2-yl}-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
 N-((5S)-3-(3-fluoro-4-(1-acetoacetyl-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
 10 oxooxazolidin-5-ylmethyl)acetamide;
 N-((5S)-3-(4-(1-(naphth-2-ylsulfonyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide;
 N-((5S)-3-(4-(1-(naphth-2-oxyacetyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide; and
 15 N-((5S)-3-(4-(1-(methylthioacetyl)-1,2,5,6-tetrahydropyrid-4-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide.

9. A process for the preparation of a compound of the formula (I) as claimed in claim 1, which comprises :-

(a) the deprotection of a compound of formula (II):


20



(II)


(b) the modification of a substituent in, or the introduction of a substituent into,
 25 another compound of formula (I) or (II);

- (c) when R^1 or R^{20} is of the formula $-NHS(O)_n(1-4C)alkyl$, wherein n is 1 or 2, the oxidation of a compound of the formula (I) or (II) wherein n is 0 or, when n is 2 the oxidation of a compound of the formula (I) or (II) wherein n is 1;
- (d) when R^1 or R^{20} is of the formula $-NHC(=O)R^b$ or $NHS(O)_n(1-4C)alkyl$, the introduction of the group $-C(=O)R^b$ or $-S(O)_n(1-4C)alkyl$ into a compound of the formula (III);




(III)

- 10 (e) when R^1 or R^{20} is hydroxy, the reaction of a compound of the formula (V) with a compound of formula (VI):




(V)




(VI)

- 15 (f) when $>A1-B1-$ is $>C=CR^{a1}-$, the reaction of a compound of the formula (VII) with a compound of the formula (VIII):




(VII)




(VIII)

- (g) when $>A1-B1-$ is $>CHCH(R^{a1})-$, the catalytic hydrogenation of a compound of the formula (I) or (II) wherein $>A1-B1-$ is $>C=CR^{a1}-$;
- (h) when $>A1-B1-$ is $>C=CR^{a1}-$, the elimination of the elements of water, or $HOCOR^{23}$ or $HOSO_2 R^{24}$ from a compound of the formula (IX) (ie. when R^{25} is $-H$, $-COR^{23}$ or $-SO_2 R^{24}$);



(IX)

- (i) when D is NR^7 and R^7 is $R^{10}CO-$ or $R^{10}S(O)_n-$, wherein n is 2, the reaction of 10 a compound of formula (X) with a compound of the formula (XI) or (XII), wherein n is 2:



(X)




(XI)



(XII)

15

- (j) when R^1 or R^{20} is azido, the reaction of a compound of the formula (XIII) with a source of azide:



(XIII)

- 5 (k) when R^1 or R^{20} is amino, the reduction of a compound of the formula (I) or (II) wherein R^1 or R^{20} is azido;
- (l) the preparation when R^1 or R^{20} is chloro, fluoro, (1-4C)alkanesulfonyloxy, or (1-4C)alkylaminocarbonyloxy, or R^{20} is of the formula $-N(CO_2R^{27})CO(1-4C)alkyl$ from a compound of the formula (I) and (II) wherein R^1 or R^{20} is hydroxy; or
- 10 (m) the preparation when R^1 or R^{20} is chloro, (1-4C)alkylthio or (1-4C)alkoxy, from a compound of the formula (XIII);
- wherein R^2 , R^3 and R^6 are as hereinabove defined; R^{18} is R^4 or protected R^4 ; R^{19} is R^5 or protected R^5 ; R^{20} is R^1 or protected R^1 ; $>A1-B1-$ is $>A-B-$ or protected $>A-B-$ and D1 is D in which functional groups are optionally protected; R^{a1} is R^a or protected R^a ;
- 15 R^{21} is (1-6C)alkyl or benzyl; R^{22} is of the formula (1-4C) alkyl or $-S(O)_n(1-4C)alkyl$; R^{23} is (1-4C)alkyl; R^{24} is an optionally substituted phenyl group; R^{25} is hydrogen, $-COR^{23}$ or $-SO_2R^{24}$; R^{26} is mesyloxy or tosyloxy; R^{27} is (1-4C)alkyl or benzyl; n is 0, 1 or 2 unless otherwise stated above; L^1 is an iodo or triflate leaving group; L^2 is a leaving group; and Y is a trialkyltin residue or a boronate acid or ester residue; and
- 20 thereafter if necessary:
- removing any protecting groups;
 - forming a pharmaceutically-acceptable salt;
 - forming an in-vivo hydrolysable ester;
- and when an optically active form of a compound of the formula (I) is required, it may
- 25 be obtained by carrying out one of the above procedures using an optically active starting material, or by resolution of a racemic form of the compound or intermediate

using a standard procedure; and similarly, when a pure regioisomer of a compound of the formula (I) is required, it may be obtained by carrying out one of the above procedures using a pure regioisomer as a starting material, or by resolution of a mixture of the regioisomers or intermediates using a standard procedure.

5 10. A pharmaceutical composition which comprises a compound of the formula (I); a pharmaceutically-acceptable salt thereof, or an in-vivo-hydrolysable ester thereof, as claimed in claims 1 to 8 and a pharmaceutically-acceptable diluent or carrier.

11. The use of a compound of the formula (I), a pharmaceutically-
10 acceptable salt thereof, or an in-vivo-hydrolysable ester thereof, as claimed in claims 1 to 8, in the manufacture of a novel medicament for use in the production of an antibacterial effect in a warm blooded animal such as man.

12. A method for producing an antibacterial effect in a warm blooded animal, such as man, in need of such treatment, which comprises administering to said animal
15 an effective amount of a compound of the present invention, a pharmaceutically-acceptable salt thereof, or an in-vivo-hydrolysable ester thereof.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 97/00462

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D413/10 A61K31/42 C07D263/24 C07D263/18 C07D417/10
 //(C07D413/10,309:00),(C07D413/10,211:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 93 09103 A (UPJOHN CO) 13 May 1993 see the whole document ---	1-12
Y	EP 0 352 781 A (DU PONT) 31 January 1990 see the whole document ---	1-12
Y	EP 0 316 594 A (DU PONT) 24 May 1989 see the whole document ---	1-12
Y	EP 0 657 440 A (HOFFMANN LA ROCHE) 14 June 1995 see the whole document ---	1-12
Y	WO 95 07271 A (UPJOHN CO ;BARBACHYN MICHAEL R (US); BRICKNER STEVEN J (US); HUTCH) 16 March 1995 see the whole document ---	1-12
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

10 June 1997

Date of mailing of the international search report

17.07.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Stellmach, J

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.

PCT/GB 97/00462

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 93 23384 A (UPJOHN CO ;HUTCHINSON DOUGLAS K (US); BRICKNER STEVEN JOSEPH (US);) 25 November 1993 see the whole document ---	1-12
Y	JOURNAL OF MEDICINAL CHEMISTRY, vol. 39, no. 3, 2 February 1996, pages 680-685, XP000574382 BARBACHYN M R ET AL: "IDENTIFICATION OF A NOVEL OXAZOLIDINONE (U-100480) WITH POTENT ANTIMYCOBACTERIAL ACTIVITY" cited in the application see the whole document ---	1-12
Y	JOURNAL OF MEDICINAL CHEMISTRY, vol. 39, no. 3, 2 February 1996, pages 673-679, XP000574381 BRICKNER S J ET AL: "SYNTHESIS AND ANTIBACTERIAL ACTIVITY OF U-100592 AND U-100766, TWO OXAZOLIDINONE ANTOBACTERIAL AGENTS FOR THE POTENTIAL TREATMENT OF MULTIDRUG-RESISTANT GRAM-POSITIVE BACTERIAL INFECTIONS" cited in the application see the whole document ---	1-12
Y	ANTIMICROB. AGENTS CHEMOTHER., vol. 31, no. 11, November 1987, pages 1791-1797, XP000654550 SLEE, A.M. ET AL.: "Oxazolidinones, a New Class of Synthetic Antibacterial Agents: In Vitro and In Vivo Activities of DuP 105 and DuP 721" see the whole document ---	1-12
Y	ANTIMICROB. AGENTS CHEMOTHER., vol. 40, no. 2, February 1996, pages 481-484, XP000654578 SPANGLER, S.K. ET AL.: "Activities of RPR 10692 (a New Oral Streptogramin), Cefditoren (a New Cephalosporin), Two New oxazolidinones (U-100592 and U-100766), and Other Oral and Parenteral Agents against 203 Penicillin-Susceptible and -Resistant Pneumococci" see the whole document ---	1-12

-/--

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/GB 97/00462

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	EXP.OPIN.INVEST.DRUGS, vol. 6, no. 2, February 1997, pages 151-158, XP000654528 ZURENKO,G.E. ET AL.: "Oxazolidinone antibacterial agents: development of the clinical candidates eperezolid and linezolid" see the whole document ---	1-12
E	WO 97 10223 A (UPJOHN CO ;GADWOOD ROBERT C (US); KAMDAR BHARAT V (US)) 20 March 1997 see the whole document -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/GB 97/00462

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9309103 A	13-05-93	AT 146783 T	15-01-97
		AU 667198 B	14-03-96
		AU 2689892 A	07-06-93
		CA 2119556 A	13-05-93
		DE 69216251 D	06-02-97
		DE 69216251 T	15-05-97
		EP 0610265 A	17-08-94
		JP 7500603 T	19-01-95
		US 5565571 A	15-10-96

EP 0352781 A	31-01-90	US 4948801 A	14-08-90
		AU 622465 B	09-04-92
		AU 3911589 A	01-02-90
		CA 1337526 A	07-11-95
		JP 2124877 A	14-05-90
		US 5130316 A	14-07-92
		US 5043443 A	27-08-91
		US 5254577 A	19-10-93

EP 0316594 A	24-05-89	AU 2404388 A	27-04-89
		CA 1317594 A	11-05-93
		DE 3884563 D	04-11-93
		DE 3884563 T	17-02-94
		ES 2059467 T	16-11-94
		JP 1135777 A	29-05-89
		SU 1801109 A	07-03-93
		US 4977173 A	11-12-90

EP 0657440 A	14-06-95	AU 8023394 A	22-06-95
		BR 9404949 A	08-08-95
		CA 2135861 A	14-06-95
		CN 1109054 A	27-09-95
		CZ 9403126 A	16-08-95
		HU 72315 A	29-04-96
		JP 7196634 A	01-08-95
		NO 944802 A	14-06-95
		NZ 270083 A	27-02-96
		PL 306244 A	26-06-95
		US 5574055 A	12-11-96
		ZA 9409710 A	13-06-95

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/GB 97/00462

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9507271 A	16-03-95	AU 7557094 A	27-03-95
		CA 2168560 A	16-03-95
		CN 1130379 A	04-09-96
		EP 0717738 A	26-06-96
		JP 9502436 T	11-03-97
		ZA 9405894 A	05-02-96

WO 9323384 A	25-11-93	AU 668733 B	16-05-96
		AU 4287793 A	13-12-93
		CN 1079964 A	29-12-93
		CZ 9402505 A	16-08-95
		EP 0640077 A	01-03-95
		FI 945246 A	08-11-94
		HU 72296 A	29-04-96
		HU 9500659 A	28-11-95
		JP 7506829 T	27-07-95
		NO 944237 A	04-01-95
		SK 133794 A	07-06-95
		US 5547950 A	20-08-96
		ZA 9302855 A	24-10-94

WO 9710223 A	20-03-97	NONE	
